

Antioxidant and antimutagenic properties of calcium sennosides in γ -Irradiated human blood cultures

M.M. Ahmed^{1*}, Z.S. Said², S.A. Montaser¹, G.A. El-Tawil¹

¹Department of Radiation Biology, National Centre for Radiation Research and Technology (NCRRT), Egyptian Atomic Energy Authority, Nasr City, Egypt

²Department of Radiation Safety, Egyptian Nuclear and Radiological Regulatory Authority (ENRRA), Nasr City, Egypt

ABSTRACT

Background: Calcium sennosides are the main active metabolites of sennas, which have a powerful interest to phytochemical and pharmacological research, due to their brilliant medicinal values. It is well known in folk medicine for their laxative and purgative uses. **Materials and Methods:** This experiment aimed to assess cytogenetic (micronucleus assay and chromosomal aberration study) and biochemical effects of calcium sennosides at a working dose (24 or 48 mg/ml) on suppressing radiation hazards in human blood cultures. Biochemical investigations include superoxide dismutase (SOD), catalase (CAT), tumor necrosis factor- α (TNF- α), interleukin-8 (IL-8), cyclooxygenase-2 (COX-2) and lactate dehydrogenase (LDH) levels. The treatment periods were 48 and 72 hrs post-irradiation at a dose of 3 Gy. Triple blood cultures for each blood sample were set up. **Results:** Ionizing radiation exposure induced significant increase in micronuclei (MN) frequencies in both mono- and bi-nucleated cells, and all types of chromosome aberrations, beside significant decrease in SOD and CAT activities. While TNF- α , IL-8, COX-2 and LDH levels were significantly increased after irradiation. Treatment with calcium sennosides exhibited decreased of micronuclei and chromosome aberration numbers and enhancement in the level of SOD and CAT activities. In addition, a significant amelioration in IL-8, TNF- α , COX-2 levels and LDH activity were scored. **Conclusion:** Present results revealed the antimutagenic and the anti-inflammatory effects of sennosides against oxidative stress induced by γ -irradiation.

Keywords: Calcium sennosides, γ -rays, pro-inflammatory cytokines, micronucleus, chromosomal aberration.

► Original article

*Corresponding authors:

Dr. A. Mahmoud Mohamed,

Fax: + 202 22748298

E-mail:

mahmoud_70mahameds@yahoo.com

Revised: June 2017

Accepted: June 2017

Int. J. Radiat. Res., July 2018;
16(3): 323-332

DOI: 10.18869/acadpub.ijrr.16.3.323

INTRODUCTION

Exposure to ionizing radiation is an integral part of our lives in different ways, such as medical, environmental, and/or even accidental. Thus, accepting of the mechanisms by which radiation toxicity develops is crucial to statement acute and chronic health problems that occurs following ionizing radiation exposure. Direct exposure to ionizing radiation-induced free radicals is totally recognized as significant contributors to early and late effects of ionizing radiation. This includes, but is not limited to, inflammation,

cytotoxicity and genotoxicity. Damage to the critical biomolecules leading to unfavorable long-term alterations in metabolic redox homeostasis following ionizing radiation exposure has been the focus of various independent investigations over last several decades ⁽¹⁾.

The first variety of senna plants were growing on the Nile River banks in Egypt and Sudan, while the second type is widely cultivated in southern and eastern parts of India. Senna plant is a small shrub classified as casual piniaceae family belongs to two genus of Cassia C senna, known as Alexandrian senna, and

Angustifolia C senna that named Tinnevely senna ⁽²⁾. Over the past 6 decades, a large number of aminothiols, natural occurring products, immunomodulators and so on were investigated for their radioprotective potential ⁽³⁾.

The biological dosimetry utilizing dicentric chromosomes examination in human lymphocytes is a well-known method accomplished since long ago together with physical dosimetry for radiation dose assessment in potentially overexposed people as well as for supposed exposures to estimate risk of health effects. The frequency of micronuclei is also commonly used as a cytogenetic biomarker. Another cytogenetic endpoint, cytokinesis-block micronucleus assay, is considered to be simple in terms of scoring criteria as a reliable and sensitive cytogenetic biomarker ⁽⁴⁾. Moreover, several researches try to manipulate different effects of sennosides such as: protective antioxidant properties, immune stimulating potency in addition to anti-inflammatory strength and its health benefits ^(5, 6).

In addition, SOD and CAT enzymes involved in oxidative stress have recently been used to monitor the development and extent of oxidative stress damage ⁽⁷⁾. On the other hand, IL-8 and TNF- α were categorized as pro-inflammatory cytokines predominantly produced by activated macrophages. Pro-inflammatory cytokines could explain most of the local and systemic components of acute and chronic inflammatory diseases. In particular IL-8 is a potent chemotactic agent for neutrophils and induces their lysosomal release ⁽⁸⁾.

COX known as prostaglandin synthase, is a membrane -bound enzyme responsible for oxidation of arachidonic acid to prostaglandins with two isoforms Cox-1 and Cox-2 had been identified. COX-1 enzyme is constitutively expressed and regulates a number of protection functions such as vascular hemostasis and gastro protection, whereas at the sites of inflammation COX-2 is inducible by number of mediators such as growth factors, cytokines and endotoxins ⁽⁹⁾. Ionizing radiation was established to induce LDH, lactate production, and extracellular acidification in human and animal

tissue in a dose dependent manner ⁽¹⁰⁾. The objective of the current study was to evaluate the radio-protective role of different doses of sennosides in irradiated human blood cultures.

MATERIALS AND METHODS

Chemicals

The chemicals for the blood cultures were purchased from GIBCO-BRL, USA. FA, cytochalasin-B, heat-inactivated foetal calf serum (FCS) and other chemicals and solvents were purchased from Sigma/ Aldrich Chemical Co., St. Louis, USA. Calcium sennosides was purchased from EL-NILE CO. for Pharmaceutical and chemical industries, Egypt, which is available in the form of tablets with the trade name (Senna Lax). Each tablet contained 12 mg of calcium sennosides. The contents of the tablet were dissolved in distilled water and withdrawn by a syringe.

Blood sampling

In the purpose of avoiding possible inter-individual variability in response to treatments, blood samples were obtained from equivalent three healthy females (average age 35 years and non-smokers) who offered an informed permission for contribution in this study. Donors were chosen carefully according to International Programme on Chemical Safety guidelines for the monitoring of genotoxic effects of carcinogens in humans ⁽¹¹⁾. Venous blood samples were collected under sanitary conditions in heparinised vacutainer test tubes (V= 5 ml, Becton Dickinson, USA) containing lithium heparin to avoid blood coagulation. The study was approved by the Central Scientific Publishing Committee, Egyptian Atomic Energy Authority, RF-156, 5-2016.

Experimental design

For each donor sample, blood was divided into 6 groups: in each group 3 samples were processed (n=3). Experimental groups were intended as the following: Group 1: control blood, groups 2 & 3: in these groups blood cultures were

treated with two doses of calcium sennosides (24 & 48 mg/ml) respectively. Group 4: blood samples were exposed to 3Gy of γ -rays. Finally 5 & 6 groups; they combined both treatment, radiation exposure (as in group 4) and calcium sinusitis treatment (as in groups 2 & 3). Two blood cultures were carried on: the first one was cultured for 48 hrs for chromosome aberration analysis and biochemical investigations. The second part was cultured for 72 hrs for micronucleus test and same biochemical parameters.

Blood culture

To 0.5 ml of the whole blood, 5ml culture medium (RPMI-1640) supplemented with 20% FCS, 200 mM l-glutamine, penicillin 100 units/ml and streptomycin 100 μ g/ml were added in 15 ml conical tubes. Phytohaemagglutinin-M with dose 0.2 ml was added to the culture to initiate cell division. Then, cells were incubated at 37°C.

Irradiation source

γ -rays were delivered to cell cultures through Canadian ^{137}Cs source that belongs to National centre of Radiation Research and Technology, Atomic Energy Authority, EGYPT. The adjusted dose was 3 Gy. The dose rate was 0.42 Gy/min. The samples were kept at 37°C immediately after irradiation till the treatment periods in the cultures with calcium sennosides started.

Cytogenetic analysis:

Cytokinesis-blocked micronucleus assay (CBMN)

CBMN was assayed by blocking cells at the cytokinesis stage by the method of Fenech ⁽¹²⁾. In each sample, a total of 1000 mono- and binucleated cells (Mono. and BN. Cells) were scored and the frequencies of cells with one, two, three micronuclei (MN) were recorded.

Chromosomal aberration analysis

The structural chromosome aberration analysis was performed according to current International Atomic Energy Authority (IAEA) guidelines ⁽¹³⁾. A total number of 100 metaphases were scored for each sample. Each

type of aberrations, as well as the percentage of aberrant cells per subject was evaluated.

Biochemical estimations

Determination of SOD activity:

In SOD activity determination Xanthine-xanthine oxidase used to generate $\text{O}_2^{\bullet-}$ and nitroblue tetrazolium (NBT) decline was used as an indicator of $\text{O}_2^{\bullet-}$ production. SOD will compete with NBT for $\text{O}_2^{\bullet-}$; the percent inhibition of NBT reduction is a measure of the amount of existing SOD. SOD activity was measured in both 48 & 72 hrs cell culture supernatants. For each tube 0.5 cacodylic buffer 0.1ml of triton X100, 0.25 ml of NBT and 1 ml of cell culture supernatants were mixed, then the mixture was incubated for 5 min at 37°C then start the reaction by adding 0.1 ml of pyrogallol, after 5 min add 0.3 ml stop solution to stop the reaction. 1.0 ml of distilled water was added instead of cell culture supernatant in blank. Finally at 540 nm optical density was measured spectrophotometrically. This quantitative method was carried on according to ⁽¹⁴⁾.

Determination of Catalase activity:

Catalase activities were measured in cell culture supernatant, according to Johansson and Borg ⁽¹⁵⁾. Reaction of catalase enzyme with methanol in the presence of an optimal concentration of hydrogen peroxide was the idea of work. 400 μ l of cell culture supernatant, 50 μ l of methanol, 50 μ l of $\text{KH}_2\text{PO}_4/\text{NaOH}$ and 10 μ l hydrogen peroxide were mixed. Mixture was incubated in shaking water bath (WB) for 20 min at 20 °C. 50 μ l of potassium hydroxide and 100 μ l of purpald solution was added to the mixture. Incubated again in WB for 10 min. at 20 °C then 50 μ l of potassium periodate was added to the mixture then centrifuged at 9500 g for 10 min. Colour compound was developed from the reaction of purpald and formaldehyde was measured spectrophotometrically at 550 nm.

Pro-inflammatory enzymes: (Cox-2 & LDH):

In the present study both COX-2 and LDH activities were measured in cell cultures supernatant via Enzyme-Linked Immunosorbent

Assay (ELISA) kit purchased from MYBioSource using double-sandwich ELISA technique. The pre-coated antibody is Human COX-2 as well as LDH monoclonal antibody and the detecting antibody is polyclonal antibody with biotin labelled. Samples and biotin labelling antibody were added into ELISA plate wells and carefully washed out with phosphate buffer saline (PBS). Then Avidin-peroxidase conjugates were added to wells in order. 3,3',5,5'-Tetramethylbenzidine (TMB) substrate was used for colouring after reactant carefully washed out by PBS. TMB changed in colour into blue in peroxidase catalytic then finally into yellow under the action of acid. The colour depth and the testing factors in samples were positively correlated.

Pro-inflammatory Cytokines : (IL-8 & TNF- α):

Human IL-8 and TNF- α were measured using solid phase sandwich ELISA from Abcam. Abcam's IL-8 as well as TNF- α in vitro Simple Step ELISA kit was used for the quantitative measurement of IL-8 & TNF- α protein in Human cell culture supernatant.

Statistical analysis

Data are presented as distribution analysis, means $\pm$ S.E. and analysed using two ways analysis of variance "F" test according to Abramowitz and Stegun ⁽¹⁶⁾. The level of statistical significance was $P < 0.05$.

RESULTS

Table 1 shows significantly increments of cells with MN frequencies (1, 2 & 3) in human lymphocytes culture exposed to 3 Gy of γ -rays, compared to control and calcium sennosides groups. Binucleated cell frequencies in irradiated group were less than that of control and calcium sennosides groups. Significant improvements occurred in all percentages of micronuclei and binucleated cells after blood culture in calcium sennosides treated media.

The most representative aberration types in chromosome aberration analysis, after irradiation is breaks, fragments and dicentrics.

figure 1 shows increments of all aberration types especially, for fragments and dicentrics, after irradiation comparing with control and calcium sennosides groups. Calcium sennosides treatments ameliorated the recorded aberration numbers.

Table 2 shows marked inhibition in both SOD & CAT activities after ionizing radiation exposure in two periods of culture (48 & 72 hrs) when compared with their corresponding control groups, while treating cultures with calcium sennosides revealed marked amelioration in both enzyme activities. SOD levels exhibited a significant increase in culture 72 hrs only when compared with that of radiation group. Also, results showed that CAT activity recorded a significant increase in both treated doses of calcium sennosides (24 & 48 mg/ml) in the two culture periods (48 & 72 hrs).

On the other hand, radiation induced significant elevation in COX-2 enzyme activities in the two cultures (48 & 72 hrs) as 1.5 folds than that recorded in control group. Calcium sennosides treatments (24 & 48 mg/ml) after irradiation tends to normalize COX-2 activities with significant change from irradiated group as illustrated in figure 2.

Figure 3 shows that LDH levels exhibited a significant increase in the radiation group in both cultures 48 & 72 hrs when compared with their corresponding control groups. A marked decrease in LDH levels were detected after treatment of irradiated cultures (48 & 72 hrs.) with calcium sennosides (24 & 48 mg/ml) than that detected in irradiated cultures.

Figure 4 shows bar charts that represents the control TNF- α in human blood culture at 48 hrs as (13%) while the irradiated group recorded an increase in TNF- α concentration about (13%) more than control while calcium sennosides (24 mg/ml) treated group recorded only (7%) increase in its level & calcium sennosides (48 mg/ml) recorded amelioration in TNF- α concentration as increased only by (2%) from control group.

Furthermore, results obtained in human blood culture at 72 hrs showed that TNF- α control levels recorded (15%). Whereas in irradiated group, TNF- α level was increased by (9%) more

than control value, while treatment with calcium sennosides at doses of 24 & 48 mg/ml showed increase in its level by 3% more than that of control group.

IL-8 levels represent 16% in culture 24 & 48 hrs respectively. In irradiated groups, IL-8

levels elevated at 5% and 8% in the culture of 48 and 72 hrs respectively, more than that recorded in control group. IL-8 level declined after treatment of irradiated blood cultures (48 and 72 hrs.) with calcium sennosides 24 & 48 mg/ml to reach levels that of control group (figure 5).

Table 1. CBMN test in human blood culture in vitro treated with sennosides (24 & 48 mg/ml) after 3Gy γ -rays exposure.

Group	Control group	Sennosides (24 mg/ml) group	Sennosides (48 mg/ml) group	Radiation group	Rad. & Sennosides (24 mg/ml) group	Rad. & Sennosides (48 mg/ml) group
Mono.cells+ 0 MN	404.3 $\pm$ 9.34	418.3 $\pm$ 6.01	417.7 $\pm$ 4.34	432.3 $\pm$ 4.34	422.0 $\pm$ 2.52	427.33 $\pm$ 3.85
Mono.cells+1MN	2.0 $\pm$ 0.58	2.0 $\pm$ 0.58	2.0 $\pm$ 0.5	22.3 $\pm$ 1.45 ^{a,b,c}	13.3 $\pm$ 1.20 ^{a,b,c,d}	14.3 $\pm$ 0.88 ^{a,b,c,d}
Mono.cells+2 MN	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	11.3 $\pm$ 0.88 ^{a,b,c}	5.3 $\pm$ 0.88 ^{a,b,c,d}	8.0 $\pm$ 0.58 ^{a,b,c,d}
Mono.cells+3 MN	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	4.3 $\pm$ 0.33 ^{a,b,c}	1.6 $\pm$ 0.88 ^{a,b,c,d}	2.0 $\pm$ 1.0 ^{a,b,c,d}
Total Mono. cells	406.3 $\pm$ 9.15	420.3 $\pm$ 5.46	419.6 $\pm$ 4.67	468.3 $\pm$ 5.79 ^{a,b,c}	442.3 $\pm$ 3.18 ^{a,b,c,d}	451.6 $\pm$ 3.18 ^{a,b,c,d}
BN + 0 MN	591.0 $\pm$ 9.08	576.7 $\pm$ 5.46	577.0 $\pm$ 4.93	526.3 $\pm$ 34.51 ^{a,b,c}	535.3 $\pm$ 3.53 ^{a,b,c}	524.0 $\pm$ 2.52 ^{a,b,c,d}
BN+ 1 MN	2.6 $\pm$ 0.33	3.0 $\pm$ 0.0	3.3 $\pm$ 0.33	27.3 $\pm$ 1.77 ^{a,b,c}	17.3 $\pm$ 1.77 ^{a,b,c,d}	18.3 $\pm$ 1.20 ^{a,b,c,d}
BN+ 2 MN	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	8.6 $\pm$ 0.88 ^{a,b,c}	4.3 $\pm$ 0.33 ^{a,b,c,d}	5.0 $\pm$ 0.58 ^{a,b,c,d}
BN+ 3 MN	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	2.7 $\pm$ 0.33 ^{a,b,c}	1.0 $\pm$ 0.58	1.67 $\pm$ 0.33 ^{a,c}
Total BN	593.7 $\pm$ 9.15	579.7 $\pm$ 5.46	580.3 $\pm$ 4.67	531.7 $\pm$ 5.7 ^{a,b,c}	557.7 $\pm$ 3.18 ^{a,b,c,d}	548.3 $\pm$ 3.18 ^{a,b,c,d}
Total No. of cells with MN	4.7 $\pm$ 0.33	5.0 $\pm$ 0.58	5.3 $\pm$ 0.67	80.7 $\pm$ 2.97 ^{a,b,c}	43.0 $\pm$ 3.52 ^{a,b,c,d}	49.3 $\pm$ 3.93 ^{a,b,c,d}
Total No. of MN	4.7 $\pm$ 0.33	5.0 $\pm$ 0.58	5.3 $\pm$ 0.67	108.7 $\pm$ 4.06 ^{a,b,c}	60.6 $\pm$ 8.18 ^{a,b,c,d}	69.7 $\pm$ 7.23 ^{a,b,c,d}

Statistical significance value ($P < 0.05$).

a, Significant when compared with control group

b, Significant when compared with sennosides 24 mg/ml group

c, Significant when compared with sennosides 48 mg/ml group

d, Significant when compared with radiation group

e, Significant when compared with Rad. & (24 mg/ml) Sennosides group




Figure 1. Effect of sennosides (24 & 48 mg/ml) treatment on the total numbers of aberration types induced by γ -rays in human blood cultures.

Statistical significance value ($P < 0.05$).

* Significant when compared with control group

** Significant when compared with radiation group

Table 2. SOD and CAT activities in human blood cultures (48 & 72 hrs) treated with sennosides (24 & 48mg/ml) after γ -rays exposure.

Groups	SOD (U/ml)		Catalase (U/ml)	
	48h	72h	48h	72h
Control group	1.06±0.09	0.9±0.09	134.0±1.52	132±0.58
Sennosides (24 mg/ml) group	1.8±0.15 ^a	1.83±0.12 ^a	126±1.73 ^a	127.3±2.73
Sennosides (48 mg/ml) group	2.17±0.07 ^a	2.27±0.09 ^{a,b}	129.3±0.88	124.3±3.39
Radiation group	0.5±0.15 ^{a,b,c}	0.47±0.09 ^{a,b,c}	100.7±1.45 ^{a,b,c}	88±1.73 ^{a,b,c}
Rad. & (24 mg/ ml) Sennosides group	0.93±0.03 ^{b,c}	1.8±0.25 ^{a,c,d}	122±0.58 ^{a,c,d}	95.3±1.45 ^{a,b,c,d}
Rad. & (48 mg/ml) Sennosides group	1.6±0.06 ^{a,c,d,e}	1.67±0.07 ^{a,c,d}	125.3±1.77 ^{a,d}	115.4±3.39 ^{a,d,e}

Statistical significance value (P < 0.05).
a, Significant when compared with control group
b, Significant when compared with sennosides 24 mg/ml group
c, Significant when compared with sennosides 48 mg/ml group
d, Significant when compared with radiation group
e, Significant when compared with Rad. & (24 mg/ ml) Sennosides group




Figure 2. Cox-2 levels in human blood cultures (48 & 72 hrs) treated with sennosides (24 & 48mg/ ml) after γ -rays exposure.
Statistical significance value (P < 0.05).
* Significant when compared with control group
** Significant when compared with radiation group




Figure 3. LDH levels in human blood cultures (48 & 72 hrs) treated with sennosides (24 & 48mg/ ml) after γ -rays exposure.
Statistical significance value (P < 0.05).
* Significant when compared with control group
** Significant when compared with radiation group




Figure 4. TNF-α level in human blood cultures treated with sennosides (24 & 48mg/ ml) after γ-rays exposure.

Statistical significance value ($P < 0.05$).
 * Significant when compared with control group
 ** Significant when compared with radiation group




Figure 5. IL-8 level in human blood cultures treated with sennosides (24 & 48mg/ ml) after γ-rays exposure.

Statistical significance value ($P < 0.05$).
 * Significant when compared with control group
 ** Significant when compared with radiation group

DISCUSSION

The world health organization (WHO) has investigated that nearly 80% of the population in the world rely on traditional medicines for meeting their health care needs (17). Deleterious effects of radiation on biological systems have been shown in different ways. Therefore, various strategies have been developed to protect biological systems by means of chemical

protection through interfering in the process of radiation damage. Since introduction of cysteine as a radioprotector in 1949, various types of natural and synthetic chemicals were tested for their radioprotective properties on various biological systems by the use of different end points (18).

Calcium sennosides has been widely used in pharmacological research and Folk medicine due to its important medicinal values as laxative

& purgative drug ⁽¹⁹⁾. Besides, they have been found to exhibit anti-inflammatory, antioxidant ⁽²⁰⁾, hypoglycemic ⁽²¹⁾, antiparasmodial, larvicidal ⁽²²⁾, antimutagenic ⁽²³⁾, and anticancer activities ⁽²⁴⁾.

Direct and indirect action are the main causes of ionizing radiation damages to DNA, where DNA interacts with the reactive free radicals like •OH, •H, and e-aq generated by radiolysis of water in direct action. While indirect action is through production of other free radicals from other molecules. By its role these reactive free radicals can be scavenged by compounds called free radical scavengers thus having the ability to provide protection against damage caused by radiation. From that point of view finding and identifying of natural effective agents which could be used for protection against radiation-induced genetic damage, especially in humans is a very vital target ⁽⁴⁾.

Leaves and fruits of *C. angustifolia* contain several laxative constituents such as anthranoids, mostly calcium sennosides A and B, aloe-emodin, emodin, and chrysophanol, fatty oils, flavonoids, polysaccharides and tannins ⁽²⁵⁾. Senna has been widely used as an analgesic, febrifuge, diuretic, hepatoprotective, vermifuge and cholagogue remedy as well as for treating amoebic dysentery, tuberculosis, gonorrhoea, dysmenorrhoea, anaemia, flu, liver and urinary tract diseases ⁽²⁶⁾.

The current study evaluates calcium sennosides antioxidant, antimutagenic, and anti-inflammatory effects on cultured human blood cells. Our results pointed to calcium sennosides treatments with both doses (24 & 48 mg/ml) reduced the numbers of radiation-induced micronuclei, chromosome aberrations (especially dicentric type) and aberrant cells (table 1, figure 1). According to Fenech ⁽¹²⁾ who stated that the "cytome" concept implies that every cell in the system studied is scored cytologically for its viability status (necrosis, apoptosis), its mitotic status (mononucleated, BN, multinucleated) and its chromosomal damage or instability status (presence of micronuclei, nucleoplasmic bridge and nucleoplasmic buds) For these reasons, it is now appropriate to refer to this technique as the

cytokinesis block MN cytome (CBMN Cyt) assay. So, the presented study used a differential count of mono- and bi-nucleated cells in 1000 to compare the mitotic and the arresting ability of each group with its treatment case. The scored data revealed a low frequency of binucleate cells in radiation group which may be attributed to radiation exposure effects on mitotic and arresting rates.

The present study observations are in agreements with the discussed data that indicated that free radicals arising during radiation-exposure can lead to DNA damage. If overproduction of these radicals occurs, oxidative damage could lead to radiation-induced cytotoxicity; chromosomal damage and micronuclei expression ⁽²⁷⁾. Some phytochemical and herbal drugs may influence this process by controlling the formation of the damaging agents in certain circumstances, while other components, such as hormones may function to enhance repair mechanism ⁽²⁸⁾.

The current study also displays the effects of γ -irradiation on human blood cell cultures (48 & 72h) which revealed significant inhibition in SOD & CAT activities compared with control group due to heavy oxidative burden induced by oxidative stress as result of free radical production. While, groups treated with calcium sennosides after irradiation recorded significant improvements in mentioned parameters values as compared with irradiated group (table 2). That observed enhancement of SOD activity recorded after calcium sinusitis treatment goes hand to hand with the results found by Shanmugasundaram ⁽⁵⁾.

Imbalance between pro-oxidant and antioxidant status in the cells following to ionizing radiation was defined by Bhosle ⁽²⁹⁾. In addition to decrease in SOD activities as a results of attacking its molecules with produced Reactive Oxygen Species (ROS) causing their denaturation and partial inactivation ⁽³⁰⁾. Also, the radioprotective effect and the antioxidative potentials of calcium sennosides have been demonstrated in several study's findings as a reduction in MN, and improvement of SOD and CAT levels ⁽⁶⁾. Moreover, the inflammatory component of most human inflammatory chronic

diseases implicates the production of pro-inflammatory cytokines. From that point of view anti TNF therapy seems promising experimentally and clinically ⁽⁸⁾.

Cell response to radiation depends on the type and dose of radiation, inherent tissue sensitivity and repair, as well as modulating intracellular factors that include position in the cell cycle, oxygen concentration, and levels of antioxidants. Intracellular oxygen determines the extent of DNA damage by X-rays and γ -rays ⁽³¹⁾. The over expression of COX-2 was recorded in the proliferative basal compartment of the stratified skin epidermis results in spontaneous hyperplasia and dysplasia in transgenic mice and furthermore a sensitization for cancer development by conferring an auto-promoted skin phenotype. In multi-stage carcinogenesis, it also becomes clear that aberrant COX-2 over expression and activity are causally involved in tumor promotion and tumor progression rather than initiation ⁽³²⁾.

Current data also show a dramatically significant increase in inflammatory mediators *in vitro* after ionizing radiation exposure as measured by COX-2 and LDH levels that may be attributed to activation of cells responsible for their production. Repeated chronic unresolved inflammation plays a definite role in the initiation of cancer and then its more developments, growth ending with metastasis ⁽³³⁾. Meanwhile, treatment of inflammation and inhibiting of accused inflammatory mediators is the point of the target. Cytokines are often involved in inflammation through their effects of antigen presentation, bone marrow differentiation, cell recruitments, cell activation and adhesion molecule expression ⁽³⁴⁾. TNF- α is considered as one of commonly used inflammatory markers.

In the present findings, ionizing radiation induce increments in both TNF- α and IL-8 *in vitro* may be attributed to the production of ROS and presence of free radicals in culture and activation of cells which responsible for their production (figures 4 and 5). Also cytokines probably play a role in perpetuating and perhaps initiating uncontrolled damage processes after irradiation exposure.

CONCLUSION

The medicinal action of senna can be attributed mainly to the anthraquinone glycosides, especially calcium sennosides. It appears that the aglycone portion is responsible for its action. The breakdown of the anthraquinone glycosides in the digestive tract can occur in one of two ways. The bowel flora can directly hydrolyze them in a similar way to that of free active aglycone. Alternatively, in the presence of bile and the sugar moiety, the free aglycone can be absorbed into the blood stream and secreted later into the colon ⁽²⁾.

Funding sources

Our research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

ACKNOWLEDGMENT

The authors greatly appreciate Dr. Mahmoud Refaat Ahmed Mohamed. Air force hospital for his help, in blood sampling.

Conflicts of interest: Declared none.

REFERENCES

1. Pathak R, Cheema, AK, Boca, SM, Krager, KJ, Hauer-Jensen, M, Nukhet Aykin-Burns, N (2015) Modulation of Radiation Response by the Tetrahydrobiopterin Pathway. *Antioxidants*, **(4)**: 68-81.
2. Balasankar D, Vanilarasu K, Selva Preetha P, Rajeswari M, Umadevi, S, Debjit Bhowmik (2013) "Senna – A Medical Miracle Plant". *Journal of Medicinal Plants Studies*, **1(3)**: 41-47.
3. Shakeri-Boroujeni A, Mozdarani H, Mahmmoudzadeh M, Faeghi F (2007) Potent radioprotective effect of herbal immunomodulator drug (IMOD) on mouse bone marrow erythrocytes as assayed with the micronucleus test. *Int J Radiat Res*, **12(3)**: 221-228.
4. Samarath RM, Samarath M, Matsumoto Y (2015) Utilization of cytogenetic biomarkers as a tool for assessment of radiation injury and evaluation of radiomodulatory effects of various medicinal plants – a review. *Drug Design Devel*

- opment and Therapy, **9**: 5355–5372.
5. Shanmugasundaram R, Kalpana Devi V, Tresina Soris P, Maruthupandian A, Mohan VR (2011) Antidiabetic, antihyperlipidaemic and antioxidant activity of *Senna auriculata* (L.) Roxb. Leaves in alloxan induced diabetic rat. *International Journal of Pharm Tech Res*, **3**(2): 747-756.
6. Jothi RS, Bharathy V, Uthayakumar F (2015) Antioxidant Potential of Aerial Part of *Senna italica* Sub Species. *micrantha* Mill. *J Pharm Sci & Res*, **7**(9): 621-625.
7. Rendic S and Guengerich FP (2012) Summary of information on the effects of ionizing and non-ionizing radiation on cytochrome P450 and other drug metabolizing enzymes and transporters. *Curr Drug Metab*, **13**(6): 787-814.
8. Nielsen OH, Ainsworth MA (2013) Tumor necrosis factor inhibitors for inflammatory bowel disease. *N Engl J Med*, **369**:754–762.
9. Kulkarni SK, Jain NK, Singh A (2000) Cyclooxygenase isoenzymes and newer therapeutic potential for selective COX-2 inhibitors. *Methods Find Exp Clin Pharmacol*, **22**(5): 291.
10. Judge JL, Owens KM, Pollock SJ, Woeller CF, Thatcher TH, Williams JP, et al. (2015) Ionizing radiation induces myofibroblast differentiation via lactate dehydrogenase. *Am J Physiol Lung Cell Mol Physiol*, **309**(8): 879-87.
11. Albertini RJ, Anderson D, Douglas GR, Hagmor L, Hemminki K, Merlo F, et al. (2000) IPCS guidelines for the monitoring of genotoxic effects of carcinogens in humans. International Programme on Chemical Safety. *Mutat. Res*, **463**: 111.
12. Fenech M (2007) Cytokinesis-block micronucleus cytome assay. *Nat Proto*, **2**, 1084.
13. IAEA (2001) Cytogenetic analysis for radiation dose assessment. In: IAEA (Eds.), International Atomic Agency Technical Report Series **405**. IAEA, Vienna, 138.
14. Sun Y, Oberley LW, Li Y (1988) A simple method for clinical assay of superoxide dismutase. *Clin Chem*, **34**:479–500.
15. Johansson LH, Borg LAH (1988) A spectrophotometric method for determination of catalase in small tissue sample *Anal Biochem*, **174**: 331-336.
16. Abramowitz M and Stegun IA (1972) Handbook of mathematical functions. Dover Publications, Inc. New York, USA.
17. Sun J, Liu BR, Hu, WJ, Yu LX, Qian XP (2007) *In vitro* anticancer activity of aqueous extracts and ethanol extracts of fifteen traditional Chinese medicines on human digestive tumor cell lines. *Phytotherapy Res*, **21**(11): 1102-1104.
18. Mahdavi M and Mozdarani H (2011) Protective effects of famotidine and vitamin C against radiation induced cellular damage in mouse spermatogenesis process. *Int J Radiat Res*, **8** (4): 223-230.
19. Hennebelle T, Weniger B, Joseph H, Sahpaz S, Bailleul F (2009) *Senna alata*. *Fitoterapia*, **80**: 385-393.
20. Yen GC and Chuang DY (2000) Antioxidant properties of water extracts from *Cassia tora* L. in relation to the degree of roasting. *J Agric Food Chem*, **48**: 2760-2765.
21. Jalalpure SS, Patil MB, Pai A, Shah BN, Salahuddin MD (2004) Antidiabetic activity of *Cassia auriculata* seeds in alloxan induced diabetic rats. *Nigerian J Nat Prod Med*, **8**: 22-23.
22. Yang YC, Lim, MY, Lee HS (2003) Emodin isolated from *Cassia obtusifolia* (Leguminosae) seed shows larvicidal activity against three mosquito species. *J Agric Food Chem*, **51**: 7629–7631.
23. Yadav AS and Jaggi S (2015) Buccal micronucleus cytome assay- a biomarker of genotoxicity. *Journal of Molecular Biomarkers & Diagnosis*, **6** (3): 236-242.
24. Prasanna R, Harish CC, Pichai R, Sakthisekaran D, Gunasekaran P (2009) Anti-cancer effect of *Cassia auriculata* leaf extract in vitro through cell cycle arrest and induction of apoptosis in human breast and larynx cancer cell lines. *Cell Biol Int*, **33**(2): 127-34.
25. Siddique I, Anis M, Aref IM (2010) In Vitro Adventitious Shoot Regeneration via Indirect Organogenesis from Petiole Explants of *Cassia angustifolia* Vahl.-a Potential Medicinal Plant. *Appl. Biochem Biotechnol*, **162**(7): 2067-2074.
26. Picon PD, Picon, RV, Costa AF, Sander GB, Amaral KM, Aboy AL, Henriques AT (2010) Randomized clinical trial of a phytotherapeutic compound containing *Pimpinella anisum*, *Foeniculum vulgare*, *Sambucus nigra*, and *Cassia angustifolia* for chronic constipation. *BMC Complement. Altern Med*, **10**: 17.
27. Halliwell B (2002) Effect of diet on cancer development: is oxidative DNA damage a biomarker? *Free Radic Biol Med*, **32**: 968.
28. Fenech M (2005) The genome health clinic and genome health nutrigenomics concepts: diagnosis and nutritional treatment of genome and epigenome damage on an individual basis. *Mutagenesis*, **20**: 255.
29. Bhosle SM, Huilgol NG, Mishra KP (2005) Enhancement of radiation-induced oxidative stress and cytotoxicity in tumor cells by ellagic acid. *Clin Chim Acta*, **359**: 89.
30. Kregel KC and Zhang HJ (2007) An integrated view of oxidative stress in aging: basic mechanisms, functional effects, and pathological considerations. *Am J Physiol Regul Integr Comp Physiol*, **292**(1): 18-36.
31. Borek C (2004) Antioxidants and Radiation Therapy. *J Nutr*, **134** (11): 3207-3209.
32. Müller-Decker K (2011) Cyclooxygenase-dependent signaling is causally linked to non-melanoma skin carcinogenesis: pharmacological, genetic, and clinical evidence. *Cancer Metastasis Reviews*, **3-4**: 343-361.
33. Zamarron BF and Chen W (2011) Dual roles of immune cells and their factors in cancer development and progression. *Int J Biol Sci*, **7**(5): 651-658.
34. Zhou X, Fragala MS, McElhaney JE, Kuchel GA (2010) Conceptual and methodological issues relevant to cytokine and inflammatory marker measurements in clinical research. *Curr Opin Clin Nutr Metab Care*, **13**(5): 541-547.