

# Effect of low dose gamma-radiation upon Newcastle disease virus antibody level in chicken

M. Vilić<sup>1\*</sup>, Ž. Gottstein<sup>2</sup>, I. Ciglar Grozdanić<sup>2</sup>, K. Matanović<sup>1</sup>, S. Miljanić<sup>3</sup>,  
H. Mazija<sup>2</sup>, P. Kraljević<sup>1</sup>

<sup>1</sup>Department of Physiology and Radiobiology, Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, 10000 Zagreb, Croatia

<sup>2</sup>Department of Poultry Diseases with Clinic, Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, 10000 Zagreb, Croatia

<sup>3</sup>Division of Materials Chemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia

**Background:** The specific antibody response against Newcastle disease virus (NDV) in the blood serum of chickens hatched from eggs exposed to low dose gamma-radiation was studied. **Materials and methods:** Two groups of eggs of commercial meat chicken lines were irradiated with the dose of 0.30 Gy <sup>60</sup>Co gamma-rays before incubation and on the 19<sup>th</sup> day of incubation, respectively. The same number of eggs unexposed to gamma-radiation served as controls. After hatching the group of chicken hatched from eggs irradiated on the 19<sup>th</sup> day of incubation was not vaccinated while the group of chicken hatched from eggs irradiated before incubation was vaccinated on the 14 day. Specific serum anti-NDV antibodies were quantified by the hemagglutination inhibition assay with 4 HA units of NDV La Sota strain. **Result:** Specific antibody titres against NDV in the blood serum of chickens hatched from eggs irradiated before incubation and vaccinated on the 14<sup>th</sup> day significantly increased on the 28<sup>th</sup> day. Specific antibody titre against NDV in the blood serum of chickens hatched from eggs irradiated on the 19<sup>th</sup> day of incubation and non-vaccinated was significantly higher on the 1<sup>st</sup> and 14<sup>th</sup> day. **Conclusion:** Acute irradiation of heavy breeding chicken eggs with the dose of 0.30 Gy <sup>60</sup>Co gamma-rays before incubation and on the 19<sup>th</sup> day of incubation could have a stimulative effect on humoral immunity in chickens. *Iran. J. Radiat. Res.*, 2009; 7 (1): 27-31

**Keywords:** Gamma-radiation, low dose effects, broiler, Newcastle disease virus, antibody.

## INTRODUCTION

It has been shown that different biological effects can be obtained by using a low dose and low dose rate of ionizing radiation such as prolongation of life span <sup>(1, 2)</sup> in parallelly with activation of immune system <sup>(3)</sup>, stimulation of growth rate and develop-

ment <sup>(4)</sup>, adaptive response and bystander effect <sup>(5, 6)</sup> and modulation of gene expression <sup>(7, 8)</sup>. Furthermore, low doses and low dose rates of ionizing radiation can stimulate humoral and cellular immunity in mice <sup>(9-13)</sup>. Previous studies regarding low dose radiation effects in chickens investigated mostly hatchability, body weight and egg fertility in commercial meat chicken lines <sup>(14-19)</sup>. Although these mentioned studies showed contradictory data, it is reported that dose of gamma radiation between 0.1 to 1 Gy, especially the dose between 0.1 and 0.5 Gy <sup>(17)</sup> or between 0.06 to 0.3 Gy, <sup>(15)</sup>, could have stimulative effects on chicken egg hatchability and chicken body weight after egg irradiation before incubation. However, to date, no information of humoral immunity is known on the chicken after irradiation of eggs by acute low dose radiation.

Antigen that has been successfully used to promote humoral immune response in chickens is Newcastle disease virus <sup>(20-22)</sup>. Although the maternal antibody in the chickens is depended on the levels in their parents <sup>(22, 23)</sup> it is known that some substance could decrease <sup>(24, 25)</sup> or enhance <sup>(26, 27)</sup> humoral immunity in chickens.

Consequently, the aim of this study was to investigate the antibody response against vaccinal NDV in blood serum of

### \*Corresponding author:

Dr. Marinko Vilić, DVM

Department of Physiology and Radiobiology, Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, HR-10000 Zagreb, Croatia

Fax: +385 12390174

E-mail: marinko.vilic@vef.hr

chickens hatched from eggs exposed to the dose of 0.30 Gy gamma-rays before and on the 19<sup>th</sup> day of incubation.

## MATERIALS AND METHODS

Investigations were carried out in two independent experiments.

### Eggs

Eggs produced by commercial meat chicken lines were irradiated three hours prior to incubation (experiment 1) and on the nineteenth day of incubation (experiment 2). In each experiment, there were included the same number of eggs unexposed to gamma-radiation and served as controls. Non-irradiated eggs were retained in the same place for a period of time equal to that required for irradiated eggs. The study was reviewed and approved by an Ethics Committee of the Faculty of Veterinary Medicine University of Zagreb.

### Irradiation and dosimetry

In both experiments eggs were irradiated with the dose of 0.30 Gy gamma radiation from panoramic <sup>60</sup>Co source (activity about 3 PBq) of the Ruder Bošković Institute, Zagreb, Croatia <sup>(28)</sup>. Dose rate was about 23.84 mGy/s, and a source axis-to-egg axis distance was 3.06 m. Dosimetric measurements were performed with an ionization chamber type 2581 and a Farmer Dosimeter type 2570 (NE Technology Limited). The dose is specified as absorbed dose to water (measured free in air).

### Incubation

All eggs of each experiment were placed in the same commercial incubator Victoria (Pavia, Italy), capacity of 22100 eggs for 19 days. Incubators had automatic control of temperature (37.8 °C), humidity (60-62 % relative humidity), and incubation rack turning (each hour). On the 19<sup>th</sup> day of incubation the eggs were transferred to hatching trays located in the same incubator.

### Chickens

After the hatch chickens were housed on the floor at a temperature appropriate to each age interval. Thus, the initial temperature was 33 °C and decreased at a rate of 2 °C per week. During the experimental period the chickens were given feed and water *ad libitum*. Both groups were kept under the same conditions.

### Vaccination schedule

Chicks were vaccinated against Newcastle disease only in the experiment 1 with a commercial Newcastle disease vaccine (PESTIKAL® La Sota SPF, VETERINA d.o.o., Zagreb, Croatia) by ocularonasal route on the 14<sup>th</sup> day.

### Serological examination

From all chickens in both experiments blood samples were collected for serological analyses from day 1 up to day 28 in weekly interval. Approximately 1 ml of blood was taken from the right jugular vein. Serum was separated by centrifugation and stored at -18 °C until use. Specific serum anti-NDV antibodies were quantitated by the hemagglutination inhibition (HI) assay as described by OIE <sup>(29)</sup>. The HI titre was defined as the reciprocal of the highest serum dilution completely inhibiting agglutination. Serum antibody titres were expressed as a log<sub>2</sub> values.

### Statistical analysis

Obtained results of antibody titre against NDV were expressed as mean value (mean ± SE) and the significance of differences between the control and irradiated groups was analysed in STATISTICA <sup>(30)</sup> using Student's t-test whereas p value <0.05 was selected to indicate significance.

## RESULTS

### Experiment 1

The results of specific HI antibody titre against NDV in the blood serum of chickens hatched from eggs irradiated before


incubation and vaccinated on the 14<sup>th</sup> day are presented in figure 1. Antibody titre against NDV was significantly increased in the blood serum of chickens hatched from irradiated eggs on the 28<sup>th</sup> day ( $p < 0.05$ ).

### Experiment 2


The results of specific HI antibody titres against NDV in the blood serum of chickens hatched from eggs irradiated on the 19<sup>th</sup> day of incubation are presented in figure 2. Antibody titre against NDV in the blood serum of chickens hatched from irradiated eggs was higher than it was in the control group during first three weeks. The difference was statistically significant on the 1<sup>st</sup> day ( $p < 0.05$ ) and the 14<sup>th</sup> day ( $p < 0.05$ ).

### DISCUSSION

Our results demonstrate that acute low-dose radiation (LDR) exposure could enhance maternal HI antibody titre against NDV in chicks as well as antibody titre synthesized in chickens after vaccination (figure 1 and 2). These results suggest that enhanced antibody titre in chicken blood serum might be due to increased antibody transport from yolk sac into embryo circulation or to the direct effect on chicken B-cell development. Namely, it is known that maternal IgG (antibodies) are transferred to offspring across the yolk sac IgG receptors without degradation<sup>(31-34)</sup>. Furthermore, HAMAL *et al.*<sup>(22)</sup> reported that level of IgG in chicken plasma directly corresponds to



**Figure 1.** Serum antibody titres to Newcastle disease virus of chickens hatched from eggs irradiated with the dose of 0.30 Gy gamma-radiation before incubation and vaccinated on the 14<sup>th</sup> day of age by oculonasal route. Results are expressed as the mean  $\pm$  standard error (SE) of 10 chickens. \*Star marks represent significant differences ( $P < 0.05$ ) between means of irradiated group and control group.



**Figure 2.** Serum antibody titres to NDV of chickens hatched from eggs irradiated with the dose of 0.30 Gy gamma-radiation on the 19<sup>th</sup> day of incubation and non-vaccinated. Results are expressed as the mean  $\pm$  standard error (SE) of 10 chickens. \*Star marks represent significant differences ( $P < 0.05$ ) between means of irradiated group and control group.

IgG level in hen plasma and is about 30% of it. Our experiment showed that acute LDR exposure on the 19<sup>th</sup> day of incubation, when the absorption of IgG is the most intensive<sup>(35)</sup>, caused significant change of antibody titre against NDV in newly hatched chicks (figure 2). However, it is still unclear what mechanisms are involved in significant increase of maternal as well as specific antibodies level to NDV in irradiated groups. Yet, we suppose the acute LDR exposure could be responsible for activation of antibody transfer over the yolk sac receptors or expression of genes for mentioned receptors. Namely, WEST *et al.*<sup>(34)</sup> firstly purified and characterized chicken yolk sac membrane IgG binding protein in 18-19 day old chicken embryos which they named FcRY receptor and sequenced its gene. Furthermore, they reported an avian protein receptor present on the chicken yolk sac membrane to be responsible for transferring of antibodies from yolk to the embryonic circulation. Since low doses of radiation can modulate protein expressions as well as new, before radiation unknown proteins in mammals<sup>(36, 37)</sup>, it is not to be excluded that dose of 0.3 Gy could not have effect upon the gene expressions of receptors. On the other side, in the experiment 1 chicken hatched from eggs irradiated before incubation and vaccinated on the 14<sup>th</sup> day after hatch had a significantly higher specific antibody titre against NDV on the 28<sup>th</sup> day compared to non-vaccinated chickens. We presume that such result could be caused by stimulating effect of acute LDR exposure on: (i) many mediators, like cytokines, which have ability to stimulate the ontogeny and activation of neonatal host defenses<sup>(38)</sup> or (ii) expression of other molecules which participate in simultaneous reactions between cells during immune response<sup>(12)</sup>.

In conclusion we want to emphasize that acute exposure of 0.30 Gy gamma-rays applied before and on the 19<sup>th</sup> day of incubation of hatching eggs seems to stimulate the humoral immunity by

enhancing transport of antibodies from eggs to embryonic circulation as well as by stimulating antibody production after vaccination.

After this, certainly, lots of questions stay open and answers to them will be results of many researches of LDR in the future on humoral immunity in birds and further extending the knowledge in the field of molecular basis of antibody and IgG transport in hens.

Our further studies are in progress to test specific anti-NDV antibodies and IgG transfer from parents to eggs and from eggs to embryonic circulation and newly hatched chicks after egg exposures by different low dose radiation in different time of incubation.

## ACKNOWLEDGMENT

*This investigation was financed by the Ministry of Science, High Educations and Sport of the Republic of Croatia. The authors wish to thank the Ministry for their financial assistance.*

## REFERENCES

1. Congdon CC (1987) A review of certain low-level ionizing radiation studies in mice and guinea pigs. *Health Phys*, **52**: 593-597.
2. Caratero A, Courtade M, Bonnet L, Planel H, Caratero C (1998) Effect of a continuous gamma irradiation at a very low dose on the life span of mice. *Gerontology*, **44**: 272-276.
3. Ina Y and Sakai K (2004) Prolongation of life span associated with immunological modification by chronic low-dose-rate irradiation in MRL-lpr/lpr mice. *Radiat Res*, **161**: 168-173.
4. Luckey TD (1982) Physiological benefits from low levels of ionizing radiation. *Health Phys*, **43**: 771-789.
5. Mothersill C and Seymour C (2004) Radiation-induced bystander effects and adaptive responses—the Yin and Yang of low dose radiobiology? *Mutat Res Fundam Mol Mech Mugag*, **568**: 121-128.
6. Streffer C (2004) Bystander effects, adaptive response and genomic instability induced by prenatal irradiation. *Mutat Res Fundam Mol Mech Mugag*, **568**: 79-87.
7. Hoshi Y, Tanooka H, Miyazaki K, Wakasugi H (1997) Induction of thioredoxin in human lymphocytes with low-dose ionizing radiation. *Biochim Biophys Acta*, **1359**: 65-70.
8. Watanabe M, Suzuki K, Kodama S (2002) Specific gene expression by extremely low-dose ionizing radiation



- which related to enhance proliferation of normal human diploid cells. *International Congress Series*, **1236**: 237-239.
9. Liu SZ, Liu WH, Sun JB (1987) Radiation hormesis: its expression in the immune system. *Health Phys*, **52**: 579-583.
  10. Safwat A (2000) The immunobiology of low-dose total-body irradiation: more questions than answers. *Radiat Res*, **153**: 599-604.
  11. Liu SZ (2003) On radiation hormesis expressed in the immune system. *Crit Rev Toxicol*, **33**: 431-441.
  12. Kojima S, Nakayama K, Ishida H (2004) Low dose gamma-rays activate immune functions via induction of glutathione and delay tumor growth. *J Radiat Res*, **45**: 33-39.
  13. Ina Y and Sakai K (2005) Activation of immunological network by chronic low-dose-rate irradiation in wild-type mouse strains: analysis of immune cell populations and surface molecules. *Int J Radiat Biol*, **81**: 721-729.
  14. Mraz FR (1971) Effect of Continuous Gamma Irradiation of Chick Embryos upon Hatchability and Growth. *Radiat Res*, **48**: 164-168.
  15. Todorov B, Tchotinski D, Cvetanov I (1986) Effect of low doses gamma-radiation upon the hatchability of eggs and live weight of the broilers hatched. Final programme and Books of abstracts of XVII<sup>th</sup> annual meeting of European Society of Nuclear Methods in Agriculture, Hannover, p. 122.
  16. Zakaria AH (1989) Effect of low Doses of Gamma Irradiation Before Incubation on Hatchability and Body Weight of Broiler Chickens Hatched Under Commercial conditions. *Poult Sci*, **68**: 1150-1152.
  17. Jilo A and Löhle K (1991) Untersuchungen über den Einfluss der gammabestrahlung Co-60 von sperma und Bruteiern verschiedener hunerrassen bzw.-genotypen auf die Brutleistungen. *Monatsh Veterinärmed*, **46**: 622-625.
  18. Zakaria AH (1991) Effect of low doses of Gamma Irradiation prior to egg Incubation on Hatchability and Body Weight of Broiler Chickens. *Br Poult Sci*, **32**: 103-107.
  19. Gerrits AR and Dijk DJ (1992) Effect of X-ray irradiation of hatching eggs on hatching time, hatchability and broiler weight. *Arch Geflügelk*, **56**: 179-181.
  20. Reynolds DL and Maraqa AD (2000) Protective immunity against Newcastle disease: The role of antibodies specific to Newcastle disease virus polypeptides. *Avian Dis*, **44**: 138-144.
  21. Seal BS, King DJ, Sellers HS (2000) The avian response to Newcastle disease virus. *Dev Comp Immunol*, **24**: 257-268.
  22. Hamal KR, Burgess SC, Pevzner IY, Erf GF (2006) Maternal Antibody Transfer from Dams to Their Egg Yolks, Egg Whites, and Chicks in Meat Lines of Chickens. *Poult Sci*, **85**: 1364-1372.
  23. Al-Natour MQ, Ward LA, Saif YM, Stewart-Brown B, Keck LD (2004) Effect of different levels of maternally derived antibodies on protection against infectious bursal disease virus. *Avian Dis*, **48**: 177-182.
  24. Arai S, Kowada T, Takehana K, Miyoshi K, Nakanishi YH, Hayashi M (1996) Apoptosis in the chicken bursa of fabricius induced by X-irradiation. *J Vet Med Sci*, **58**: 1001-1006.
  25. Singhal LK, Bagga S, Kumar R, Chauhan RS (2003) Down regulation of humoral immunity in chickens due to carbendazim. *Toxicol In Vitro*, **17**: 687-92.
  26. Amer S, Na KJ, El-Abasy M, Motobu M, Koyama Y, Koge K, Hirota Y (2004) Immunostimulating effects of sugar cane extract on X-ray radiation induced immunosuppression in the chicken. *Int J Immunopharmac*, **4**: 71-77.
  27. Kong X, Hu Y, Rui R, Wang D, Li X (2004) Effects of Chinese herbal medicinal ingredients on peripheral lymphocyte proliferation and serum antibody titer after vaccination in chicken. *Int J Immunopharmac*, **4**: 975-982.
  28. Miljanić S and Ranogajec-Komor M (1997) Application of cavity theory to the response of various TLDs to <sup>60</sup>Co gammas degraded in water. *Phys Med Biol*, **42**: 1-15.
  29. OIE (2005) World Organisation for Animal Health. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals (chapter 2.1.15 on Newcastle disease). Available online at: [http://www.oie.int/Eng/Normes/Mmanual/A\\_00038.htm](http://www.oie.int/Eng/Normes/Mmanual/A_00038.htm)
  30. StatSoft Inc. (2005) Statistica (data analysis software system). Version 7.1.
  31. Kramer TT and Cho HC (1970) Transfer of immunoglobulins and antibodies in the hen's egg. *Immunology*, **19**: 157-167.
  32. Linden CD and Roth TF (1978) IgG receptors on foetal chick yolk sac. *J Cell Sci*, **33**: 317-328.
  33. Tressler RL and Roth TF (1987) IgG receptors on the embryonic chick yolk sac. *J Biol Chem*, **262**: 15406-15412.
  34. West AP Jr, Herr AB, Bjorkman PJ (2004) The chicken yolk sac IgY receptor, a functional equivalent of the mammalian MHC-related Fc receptor, is a phospholipase A2 receptor homolog. *Immunity*, **20**: 601-610.
  35. Kowalczyk K, Daiss J, Halpern J, Roth TF (1985) Quantitation of maternal-fetal IgG transport in the chicken. *Immunology*, **54**: 755-62.
  36. Chen S, Cai L, Li X, Liu S (1999) Low-dose whole-body irradiation induces alteration of protein expression in mouse splenocytes. *Toxicol Lett*, **105**: 141-152.
  37. Chen SL, Cai L, Meng QY, Xu S, Wan H, Liu SZ (2000) Low-Dose Whole-Body Irradiation (LD-WBI) Changes Protein Expression of Mouse Thymocytes: Effect of a LD-WBI-Enhanced protein RIP10 On Cell Proliferation and Spontaneous or Radiation-Induced Thymocyte Apoptosis. *Toxicol Sci*, **55**: 97-106.
  38. Kougut MH (2000) Cytokines and prevention of infectious diseases in poultry: a review. *Avian Pathol*, **29**: 395-404.

