

# Radioactivity of soil at proposed Rooppur Nuclear Power Plant site in Bangladesh

J. Ferdous\*, A. Begum, A. Islam

Health Physics Division, Atomic Energy Centre, Dhaka, Bangladesh

## ABSTRACT

### ► Original article

**\* Corresponding author:**

Dr. Jannatul Ferdous,

E-mail: ferdous28@yahoo.com

Revised: Dec. 2014

Accepted: Jan. 2015

Int. J. Radiat. Res., April 2015;  
13(2): 135-142

DOI: 10.7508/ijrr.2015.02.003

**Background:** The concentrations and distributions of natural and anthropogenic radioactive materials in soils of the site of Rooppur Nuclear Power Plant (RNPP) were investigated with the aim of evaluating the environmental radioactivity and radiation hazard. **Materials and Methods:** Soil samples were collected from 40 locations in and around the proposed site of Rooppur Nuclear Power Plant and determined the activity level by using a gamma-ray spectrometry. **Results:** The concentrations of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in soil samples, radium equivalent activity ( $\text{Ra}_{\text{eq}}$ ) for soil, absorbed dose rates, external hazard ( $\text{H}_{\text{ex}}$ ) values were determined. No artificial radioactivity ( $^{137}\text{Cs}$ ) was found in these samples. **Conclusion:** This present study provides a baseline of radioactivity and radiation levels in the proposed nuclear power plant site area at Rooppur. It will be utilized to correlate the radioactivity measured after operation of the reactor.

**Keywords:** Gamma spectrometry, radioactivity, soil, Rooppur nuclear power plant (RNPP).

## INTRODUCTION

In modern times, economic growth of a country is measured in terms of its capacity for production of electricity. All countries seek to ensure a supply of electricity that is affordable, reliable, and secure for sustaining modern lives. In developing countries like Bangladesh the demand for electricity is growing and the actual production of electricity is expanding day-by-day. In order to improve the electric power generation capacity and sustain their economic growth, long-term plan is needed. From this view, the government of Bangladesh has attached firm commitment to implement the Rooppur NPP as the first Nuclear Power Plant (NPP) in Bangladesh. Bangladesh has domestically and internationally committed to the peaceful applications of ionizing radiation in order to achieve socio-economic development in the country. Environmental radiation and radioactivity base line study has become important nationally in every country. Such

investigations can be useful for both the assessment of public dose rates and the performance of epidemiological studies, as well as to keep reference-data records, to ascertain possible changes in the environmental radioactivity due to nuclear, industrial and other human activities.

Radiation is a natural part of the environment in which we live (1-4). All people receive exposure from natural origin (e.g.,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{226}\text{Ra}$ ,  $^{222}\text{Rn}$ ,  $^{210}\text{Pb}$ ,  $^{40}\text{K}$  etc.), as well as anthropogenic ones ( $^{137}\text{Cs}$ ,  $^{134}\text{Cs}$ ,  $^{131}\text{I}$ ,  $^{90}\text{Sr}$ ,  $^{85}\text{Kr}$ ,  $^{239}\text{Pu}$  etc.) radionuclides in soil, water, air and food. The natural occurrence of  $^{238}\text{U}$  and  $^{232}\text{Th}$  in the earth's crust is a source of external exposure due to gamma-rays. The  $^{238}\text{U}$  decay series consists of 15 members in total including  $^{226}\text{Ra}$  and the series ends with stable  $^{206}\text{Pb}$  which is a stable element. After  $^{226}\text{Ra}$ , the series emits eight alpha and six beta particles along with many gamma-rays. There are 12 members in the decay series of  $^{232}\text{Th}$  and it decays into  $^{208}\text{Pb}$  after emitting seven alpha and five beta particles

along with many gamma-rays. In addition, among the naturally occurring potassium isotopes,  $^{40}\text{K}$  is unstable, and decays by the emission of 1,460 keV gamma-rays and is also a source of gamma-rays.  $^{137}\text{Cs}$  in the environment comes from a variety of sources. The largest single source is a from atmospheric nuclear weapons tests in the 1950s and 1960s, which dispersed and deposited  $^{137}\text{Cs}$  world-wide. However, much of the  $^{137}\text{Cs}$  from the testing has now been decayed. Nuclear reactor accident, such the Chernobyl accident (1986) in the Ukraine, releases  $^{137}\text{Cs}$  to the environment (5,6). Radioactivity in soil and water system may ultimately find its way to human through food chain and by direct content with the ecosystem. So, the knowledge of the distribution pathways of both artificial and natural radionuclides is essential in maintaining some surveys of control of prevailing radiation and radioactive contamination levels (7).

Bangladesh Atomic Energy Commission (BAEC) has taken a comprehensive programme on measurement of radiation level at the site of the RNPP. There are no other nuclear/radioactive installations around the NPP site. The installation of radioactivity monitoring data will be utilized to compare pre and post installation data and to take safety measures if there is any abnormal rise of radiation level due to operation power reactors.

Thus this will ensure safety of the population and the environment. The aim of this study is to determine natural ( $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ) and artificial ( $^{137}\text{Cs}$ ) radioactivity levels in soils from various areas of RNPP site, and also to investigate the radium equivalent activity ( $Ra_{eq}$ ), the absorbed dose rate, and external and internal radiation hazard indices. The result of the monitoring study will be required as regulatory pre-requisites for the installation /operation of RNPP in future.

## MATERIALS AND METHODS

### Sample collection

The NPP site is located on the east bank of the Ganges River at the village of Rooppur, the district of Pabna and approximately 170 km northwest of the city Dhaka, the capital of Bangladesh. The coordinates of the sites are latitude  $24^{\circ}4.35''$  North and longitude  $89^{\circ}2.80''$  East. Soil samples were collected from 40 locations around the proposed site of RNPP. The locations are shown in the figure 1. All the samples were collected at 0-5 cm depth from the soil-surface. Each of the samples weighed 1kg. The samples transferred in dried acetone-cleaned polyethylene bags with sample codes were transported to the laboratory.




Figure 1. Location Map of Rooppur (Bangladesh) Nuclear Power Plant site.

### Sample preparation

At the lab, the collected samples were transferred from the polyethylene bags to the acetone-cleaned stainless steel bucket and dried in an oven at 100-105 °C until a constant weight was achieved. Each of the dried samples was ground to fine powder in an agate motor separately. The powdered soil samples were then sieved using a fine aperture mesh screen (mesh size 2 mm) in order to remove extraneous items like plant materials, roots, pebbles etc. and to obtain a fine-grained sample that would present a uniform matrix to the detector. The ground samples were stored in separate polyethylene packets as stock samples. Finally, each of the samples was transferred to cylindrical plastic-container. The containers were approximately of equal size and shape (i.e., diameter 6.5 cm and height 7.5 cm). The net weight of each sample was determined using a microbalance. The weight is different for each sample because of the different density of the sample material. The containers were then sealed tightly, rapped with thick vinyl tapes around their screw necks. The samples were stored for at least four weeks to reach secular equilibrium between the  $^{238}\text{U}$  and  $^{232}\text{Th}$  series and their respective progenies to get ready for measurements (8).

### Measurement procedures

Since 98.5% of the radiological hazard of the Uranium series are caused by the Radium, and its daughter products and the contribution from the  $^{238}\text{U}$  and the other daughter products of  $^{226}\text{Ra}$  can generally be ignored (9). The radioactivities of soil samples were measured using a low-level gamma counting system, a high resolution HPGe coaxial detector (EG & G ORTEC) coupled with a silena Emcaplus multichannel analyzer (MCA) and associated microprocessors. The effective volume of the detector was 83.469 cm<sup>3</sup> and energy resolution of the 1.33 MeV energy peak for  $^{60}\text{Co}$  was found as 1.69 keV at full width half maximum (FWHM) with a relative efficiency of 19.6%. The detector was shielded by using requisite thickness of mild steel ring at the side and lead at the top for optimization of the background radiation level.

The radioactivity concentration of  $^{226}\text{Ra}$  was determined from  $\gamma$ -ray energies of its daughters  $^{214}\text{Pb}$  (351.92 & 295.21 keV) and  $^{214}\text{Bi}$  (609.31 & 1120.30 keV), while the  $^{232}\text{Th}$  was determined from  $\gamma$ -ray energies of its daughters  $^{212}\text{Pb}$  (238.63 keV),  $^{208}\text{Tl}$  (583.14 keV) and  $^{228}\text{Ac}$  (911.07 & 969.11 keV). The radioactivity concentrations of  $^{40}\text{K}$  and  $^{137}\text{Cs}$  were determined from their  $\gamma$ -ray energy of 1460.80 keV and 662 keV respectively. The efficiency calibration of the detector was performed by using standard sources Eu-152 mixed with 250 g of analytical grade  $\text{Al}_2\text{O}_3$  at different energy such as 121.78 keV, 244.69keV, 344.27 keV, 411.11 keV, 443.91 keV, 778.89 keV, 963.38 keV, 1085.78 keV, 1112.01 keV & 1407.67 keV. The efficiency calibration curve thus obtained was employed in the different energy peaks covering the range up to 2000 keV to obtain the efficiency of the detector for the particular gamma ray energy of interest. The geometry of the counting samples was the same as that of the standard samples and the counting time for all the sample was 5000sec. In order to determine the background distribution in the environment around the detector, an empty sealed beaker was counted in the same manner and in the same geometry as the samples. The background spectra were used to correct the net peak area of gamma rays of measured isotopes.

### Theoretical calculations

*The activity concentrations:* The activity concentrations of the radionuclides in the measured samples were computed using the following relation (10),

$$A_s (\text{BqKg}^{-1}) = \frac{C_a}{\epsilon P_\gamma M_s} \quad (1)$$

where  $C_a$  is the net gamma counting rate (counts per second),  $\epsilon$  the detector efficiency of the specific  $\gamma$ -ray,  $P_\gamma$  the transition probability of gamma decay and  $M_s$  is the mass of the sample (kg).

### Radium equivalent activity

For the purpose of comparing the radiological effect or activity of materials that contain  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  by a single quantity,

which takes into account the radiation hazards due to these isotopes, a common index termed as the radium equivalent activity ( $Ra_{eq}$ ) is used. This activity index provides a useful guideline in regulating the safety standards on radiation protection for the general public residing in the area under investigation. The  $Ra_{eq}$  index represents a weighted sum of activities of the above mentioned natural radionuclides. The index is given as <sup>(3)</sup>:

$$Ra_{eq} = A_{Ra} + 1.43A_{Th} + 0.077A_K \quad (2)$$

where  $A_{Ra}$ ,  $A_{Th}$  and  $A_K$  are the average activity concentration in the sample in  $Bqkg^{-1}$  of  $^{226}Ra$ ,  $^{232}Th$  and  $^{40}K$  respectively.

### The absorbed dose rate

The absorbed dose rate in air at average gonad height of one meter above the surface of ground due to the natural radionuclides  $^{226}Ra$ ,  $^{232}Th$  and  $^{40}K$  was estimated using formula given in UNSCEAR, 1988<sup>(3)</sup>:

$$D = [0.427A_{Ra} + 0.662A_{Th} + 0.0432A_K] nGyh^{-1} \quad (3)$$

where  $A_{Ra}$ ,  $A_{Th}$  and  $A_K$  are the same meaning as in equation 2.

### External and Internal Hazard indices

The soils are used for manufacturing earthen huts, bricks and pottery materials and, hence, the external radiation hazard index,  $H_{ex}$  and internal radiation hazard index,  $H_{in}$  were calculated using the following relations <sup>(8)</sup>.

$$H_{ex} = \frac{A_{Ra}}{370} + \frac{A_{Th}}{259} + \frac{A_K}{4810} \leq 1 \quad (4)$$

$$H_{in} = \frac{A_{Ra}}{185} + \frac{A_{Th}}{259} + \frac{A_K}{4810} \leq 1 \quad (5)$$

where  $A_{Ra}$ ,  $A_{Th}$  and  $A_K$  are the same meaning as in equation 2.

## RESULTS AND DISCUSSION

### Activity concentrations of Natural Radionuclides

The calculated activity concentrations of radionuclides were obtained from gamma spectrometry measurements for 40 soil samples collected from different locations of the Rooppur

NPP site, Bangladesh in table 1. Also shown are the derived radium equivalent activity, the absorbed dose rate, and external and internal radiation hazard indices due to activity concentration of the natural radionuclides measured. Three major natural gamma radiation sources were identified. Radium-226 was the precursor of most of the  $\gamma$ -emitting radionuclide identified in the Uranium decay series. Thorium-232 was likely the precursor of the isotopes identified in the Thorium decay series. Potassium-40 which accounts for most of the activity is the third source <sup>(9)</sup>. It should be noted that the values given in the tables 2 and 3 are not the representative values for each country but for the locations from where the samples have been collected. For  $^{226}Ra$  activity concentration, each value was obtained from the 351.92 keV photo-peaks of  $^{214}Pb$  and the 609.31 keV photo-peaks of  $^{214}Bi$ . The results for soil ranged from  $25.52 \pm 1.37$  to  $37.54 \pm 2.28 Bqkg^{-1}$  with an average of  $30.85 \pm 3.42 Bqkg^{-1}$ . For the  $^{232}Th$  activity concentration, each value was an average of three values obtained from 238.63keV photo-peaks of  $^{212}Pb$ , 911.07 and 969.11 keV photo peaks of  $^{228}Ac$ . The results for soil ranged from  $32.89 \pm 1.66$  to  $53.37 \pm 2.00 Bqkg^{-1}$  with an average of  $40.88 \pm 5.17 Bqkg^{-1}$ . The activity concentration of  $^{40}K$  was obtained from its photo-peak of 1460.80 keV. The results for soil ranged from  $313.19 \pm 27.76$  to  $521.65 \pm 28.35 Bqkg^{-1}$  with the average value of  $390.10 \pm 53.84 Bqkg^{-1}$ . The errors quoted represent the counting error for individual measurements and standard deviation for means. In the soil, in general, the activity concentration of  $^{232}Th$  was found to be higher than that of  $^{226}Ra$ , while that of  $^{40}K$  markedly exceeds the values of both  $^{226}Ra$  and  $^{232}Th$ . Table 1 shows that the activity concentration of thorium is higher than radium, which is evident from the fact that the activity of thorium is 1.5 times higher than that of Uranium in earth's crust <sup>(11)</sup>. It is also observed that the measured activity concentration of  $^{40}K$  exceeds markedly the values of both Radium and Thorium, as it is the most abundant radioactive element under consideration. Moreover the excessive use of the Potassium containing



fertilizers in the area adjacent to the sampling sites may contribute to the higher values of <sup>40</sup>K activity. In the present study, the activity concentrations of <sup>226</sup>Ra and <sup>232</sup>Th in soil are comparable with the reported values in table 3, especially those at the different regions of Bangladesh (12-15). The activity concentrations of <sup>40</sup>K are comparable with the other values

reported in table 2. Moreover, our obtained values fall within the range of corresponding world values and other published results mentioned in table 2. The world average activity concentration (2,3,8) of <sup>226</sup>Ra is 40 Bqkg<sup>-1</sup> with ranges of 15 – 50 Bqkg<sup>-1</sup>, <sup>232</sup>Th is 40 Bqkg<sup>-1</sup> with ranges of 7 – 50 Bqkg<sup>-1</sup> and <sup>40</sup>K is 580 Bqkg<sup>-1</sup> with ranges of 100 – 700 Bqkg<sup>-1</sup>.

**Table1.** Radioactivity concentrations of radionuclides as well as the calculated Ra equivalent activity, absorbed dose rate, external hazard index and internal hazard index in soil samples across the site of RNPP.

| No of Soil sample | Activity of Radionuclide (Bqkg <sup>-1</sup> ) |                   |                 |                   | Ra <sub>eq</sub> (Bqkg <sup>-1</sup> ) | Dose Rate D (nGyh <sup>-1</sup> ) | H <sub>ex</sub> | H <sub>in</sub> |
|-------------------|--|-------------------|-----------------|-------------------|--|-----------------------------------|-----------------|-----------------|
|                   | <sup>226</sup> Ra                              | <sup>232</sup> Th | <sup>40</sup> K | <sup>137</sup> Cs |  |                                   |                 |                 |
| RNPP-01           | 27.92±1.01                                     | 36.79±2.15        | 374.90±27.95    | -                 | 109.32                                 | 52.47                             | 0.295           | 0.371           |
| RNPP-02           | 30.14±2.24                                     | 36.56±2.17        | 330.59±32.93    | -                 | 107.80                                 | 51.35                             | 0.291           | 0.373           |
| RNPP-03           | 35.83±1.60                                     | 43.16±1.38        | 335.54±48.33    | -                 | 123.30                                 | 58.37                             | 0.333           | 0.430           |
| RNPP-04           | 31.71±2.10                                     | 39.73±2.15        | 396.93±30.88    | -                 | 119.00                                 | 56.99                             | 0.322           | 0.407           |
| RNPP-05           | 32.82±1.49                                     | 49.83±1.01        | 415.30±40.58    | -                 | 135.95                                 | 64.94                             | 0.367           | 0.456           |
| RNPP-06           | 29.05±1.31                                     | 38.86±1.46        | 374.39±30.08    | -                 | 113.36                                 | 54.30                             | 0.311           | 0.38            |
| RNPP-07           | 36.86±2.16                                     | 42.75±2.22        | 432.25±30.27    | -                 | 131.18                                 | 62.71                             | 0.355           | 0.454           |
| RNPP-08           | 28.05±1.59                                     | 35.98±1.29        | 376.12±33.35    | -                 | 108.38                                 | 52.04                             | 0.293           | 0.369           |
| RNPP-09           | 26.22±1.54                                     | 32.89±1.66        | 335.54±38.33    | -                 | 99.02                                  | 47.46                             | 0.268           | 0.338           |
| RNPP-10           | 34.41±1.45                                     | 46.13±2.29        | 394.45±32.36    | -                 | 130.65                                 | 62.27                             | 0.353           | 0.446           |
| RNPP-11           | 33.55±1.66                                     | 42.76±1.19        | 370.48±36.40    | -                 | 123.13                                 | 58.64                             | 0.333           | 0.423           |
| RNPP-12           | 36.91±1.77                                     | 44.32±1.45        | 354.14±38.57    | -                 | 127.466                                | 60.40                             | 0.345           | 0.444           |
| RNPP-13           | 27.86±2.11                                     | 36.43±1.88        | 322.91±31.19    | -                 | 104.74                                 | 49.96                             | 0.283           | 0.358           |
| RNPP-14           | 29.87±2.19                                     | 43.14±1.95        | 401.07±29.68    | -                 | 122.35                                 | 58.64                             | 0.331           | 0.411           |
| RNPP-15           | 30.76±2.82                                     | 40.95±2.04        | 335.05±28.67    | -                 | 113.59                                 | 54.05                             | 0.307           | 0.390           |
| RNPP-16           | 30.13±2.62                                     | 39.94±1.31        | 464.88±38.14    | -                 | 122.95                                 | 59.39                             | 0.332           | 0.414           |
| RNPP-17           | 32.11±1.72                                     | 44.04±1.99        | 409.92±39.28    | -                 | 126.57                                 | 60.57                             | 0.342           | 0.429           |
| RNPP-18           | 25.52±1.37                                     | 36.34±1.84        | 389.29±29.55    | -                 | 107.38                                 | 51.77                             | 0.291           | 0.359           |
| RNPP-19           | 39.44±2.01                                     | 50.80±2.28        | 521.65±28.35    | -                 | 152.14                                 | 73.01                             | 0.411           | 0.518           |
| RNPP-20           | 29.32±1.69                                     | 40.89±1.44        | 399.10±32.55    | -                 | 118.43                                 | 56.83                             | 0.320           | 0.399           |
| RNPP-21           | 31.00±1.59                                     | 49.37±1.34        | 381.83±30.17    | -                 | 130.90                                 | 62.41                             | 0.354           | 0.438           |
| RNPP-22           | 30.74±2.64                                     | 40.31±2.10        | 361.28±30.39    | -                 | 116.15                                 | 55.42                             | 0.314           | 0.397           |
| RNPP-23           | 29.74±1.75                                     | 41.80±2.49        | 338.40±26.25    | -                 | 115.49                                 | 54.99                             | 0.312           | 0.393           |
| RNPP-24           | 32.29±1.41                                     | 49.35±1.27        | 396.70±35.36    | -                 | 133.31                                 | 63.60                             | 0.360           | 0.448           |
| RNPP-25           | 30.47±2.02                                     | 43.85±1.39        | 402.18±32.13    | -                 | 124.05                                 | 59.41                             | 0.335           | 0.418           |
| RNPP-26           | 28.87±1.76                                     | 34.56±1.93        | 313.19±27.76    | -                 | 102.33                                 | 48.74                             | 0.277           | 0.355           |
| RNPP-27           | 27.06±1.39                                     | 38.64±2.18        | 352.53±32.73    | -                 | 109.38                                 | 52.36                             | 0.296           | 0.369           |
| RNPP-28           | 29.24±1.94                                     | 36.33±1.56        | 415.30±30.58    | -                 | 113.09                                 | 54.48                             | 0.306           | 0.385           |
| RNPP-29           | 31.52±1.61                                     | 49.97±1.79        | 383.82±27.41    | -                 | 132.43                                 | 63.12                             | 0.358           | 0.443           |
| RNPP-30           | 26.99±2.01                                     | 33.45±2.07        | 396.83±29.26    | -                 | 105.30                                 | 50.81                             | 0.286           | 0.358           |
| RNPP-31           | 28.37±1.88                                     | 40.50±1.21        | 332.76±27.99    | -                 | 111.82                                 | 53.30                             | 0.302           | 0.379           |
| RNPP-32           | 31.26±1.51                                     | 42.74±3.55        | 436.49±30.10    | -                 | 125.89                                 | 60.50                             | 0.340           | 0.425           |
| RNPP-33           | 27.04±1.74                                     | 50.18±2.21        | 518.30±28.47    | -                 | 138.59                                 | 67.16                             | 0.375           | 0.448           |
| RNPP-34           | 33.76±2.10                                     | 46.62±1.95        | 438.30±21.29    | -                 | 134.08                                 | 64.21                             | 0.362           | 0.454           |
| RNPP-35           | 37.54±2.28                                     | 53.37±2.00        | 504.90±28.31    | -                 | 152.62                                 | 73.17                             | 0.412           | 0.514           |
| RNPP-36           | 30.26±1.79                                     | 45.17±2.66        | 494.85±28.19    | -                 | 132.85                                 | 64.20                             | 0.359           | 0.441           |
| RNPP-37           | 26.54±1.99                                     | 40.07±3.52        | 361.34±27.79    | -                 | 111.58                                 | 53.47                             | 0.302           | 0.373           |
| RNPP-38           | 33.92±1.75                                     | 38.12±2.05        | 350.66±30.26    | -                 | 115.35                                 | 54.86                             | 0.311           | 0.403           |
| RNPP-39           | 25.63±1.22                                     | 39.82±2.02        | 368.04±27.87    | -                 | 110.82                                 | 53.20                             | 0.299           | 0.368           |
| RNPP-40           | 33.25±1.67                                     | 38.79±2.48        | 321.65±28.51    | -                 | 113.40                                 | 53.77                             | 0.306           | 0.396           |
| Average           | 30.85  | 40.88             | 390.10          | -                 | 120.65                                 | 57.73                             | 0.326           | 0.409           |
| STD               | 3.42   | 5.17              | 53.84           | -                 | 12.66                                  | 6.10                              | 0.034           | 0.042           |

(± values represent counting error for individual measurement and standard deviation for average. The maximum and minimum values are given in bold face).

**The fission product (<sup>137</sup>Cs)**

The manmade fission product residue <sup>137</sup>Cs, which would come down to this part of the earth from the atmosphere following the nuclear power plant accident at Chernobyl on 26 April 1986 and other previous test of nuclear devices around the world, was considered in this study to obtain an estimate of fallout in the Rooppur NPP site. But no detectable activity of <sup>137</sup>Cs was observed in the soil samples under study. It implies, therefore, that there is no activity due to fallout in the site of the Rooppur NPP, Bangladesh.

**Radium equivalent activity, Ra<sub>eq</sub>**

Calculated data of Ra<sub>eq</sub> is presented in column 6 of table 1. The Ra<sub>eq</sub> for soil varied in the range 99.02 to 152.62 Bqkg<sup>-1</sup> with the average value of 120.65±12.66 Bqkg<sup>-1</sup>. Those are far below the allowable limit (370 Bqkg<sup>-1</sup>) as recommended by the IAEA<sup>(2,3,8)</sup>.

**The absorbed dose rate**

In the present study the dose rate due to <sup>226</sup>Ra, <sup>232</sup>Th, and <sup>40</sup>K in soil samples varied from 47.46 to 73.17 nGyh<sup>-1</sup> with an average of 57.73±6.10 nGyh<sup>-1</sup>. These values are comparable with the world average <sup>(3)</sup> value of 55 nGyh<sup>-1</sup>.

**External and internal hazard index**

The external hazard index (H<sub>ex</sub>) due to natural gamma radiation was calculated using the equation (3) which is shown in column 8 of table 1. The H<sub>ex</sub> value for soil in the area of interest ranged from 0.268 to 0.412 with the average value of 0.326±0.034. The calculated H<sub>ex</sub> values for all samples should be lower than unity, which do not cause any harm to the population in all regions under investigation. All values of H<sub>ex</sub> in the present work are less than unity. There is also a radiation hazard threat to respiratory organs due to the <sup>222</sup>Rn, decay product of <sup>226</sup>Ra, and its short-lived decay products. To account for this the maximum permissible concentration for radium must be reduced to half of the normal limit <sup>(8)</sup>. Considering this limit, the internal hazard index, H<sub>in</sub> was calculated and is shown in the last column of table 1. The H<sub>in</sub> for soil is found to range from 0.338 to 0.514 with the average value of 0.409±0.042.

Table 2 and table 3 show the comparative study of the present work with others of the world. However, slight variation in the radioactivity content in soil was observed with various locations worldwide mainly due to soil type, formation and transport process involved.

Table 2. Comparison of present study with other work.

| Location  | Activity in Bqkg <sup>-1</sup> |                   |                 |                   |
|---|--------------------------------|-------------------|-----------------|-------------------|
|   | <sup>226</sup> Ra              | <sup>232</sup> Th | <sup>40</sup> K | <sup>137</sup> Cs |
| Dhaka, Bangladesh <sup>(23)</sup>                   | 37.8                           | 58.2              | 790.8           | -                 |
| Chittagong, Bangladesh <sup>(12)</sup>              | 34.6                           | 60                | 438             | 1.08              |
| Jessore, Bangladesh <sup>(22)</sup>                 | 48                             | 53                | 481             | -                 |
| Nine southern districts, Bangladesh <sup>(14)</sup> | 42                             | 81                | 833             | 2.08              |
| Eastern sichuan province, China <sup>(20)</sup>     | 26                             | 49                | 440             | 6                 |
| Peshwar, Pakistan <sup>(18)</sup>                   | 65                             | 84                | 646             | -                 |
| Nigeria <sup>(17)</sup>                             | 8.3                            | 34.3              | 684             | -                 |
| Nile Delta, Egypt <sup>(19)</sup>                   | 17                             | -                 | 316             | -                 |
| Louisiana, USA <sup>(9)</sup>                       | 43 - 95                        | 50 - 190          | 43 - 729        | 5-58              |
| Worldwide average <sup>(1)</sup>                    | 40(15-50)                      | 40(7-50)          | 580(100-700)    | -                 |
| RNPP site (Bangladesh) (Present study)              | 30.85                          | 40.88             | 390.10          | -                 |

Table 3. Comparative study of the present works with others of the world.

| Country                                    | Outdoor average absorbed dose rate in nGy h <sup>-1</sup> | Range  | Outdoor effective dose equivalent at 1.0m gonad height in μSvy <sup>-1</sup> | Method of analysis                        |
|--|---|--------|--|---|
| Romania <sup>(3)</sup>                     | 81  | 32-210 | 99.34  | Analysis of soil using gamma Spectrometry |
| Tripoli <sup>(16)</sup>                    | 23  | -      | 28.21  | Analysis of soil using gamma Spectrometry |
| Nigeria <sup>(17)</sup>                    | 128   | 5-186  | 156.90   | Analysis of soil using gamma Spectrometry |
| Pakistan <sup>(18)</sup>                   | 49  | 39-63  | 60   | Analysis of soil using gamma Spectrometry |
| Egypt <sup>(19)</sup>                      | 93  | -      | 114.6  | Analysis of soil using gamma Spectrometry |
| China <sup>(20)</sup>                      | 60  | 50-72  | -  | Analysis of soil using gamma Spectrometry |
| Six districts, Bangladesh <sup>(21)</sup>  | 77  | 59-103 | 94.93  | Analysis of soil using gamma Spectrometry |
| Nine districts, Bangladesh <sup>(14)</sup> | 107   | 60-156 | -  | Analysis of soil using gamma Spectrometry |
| Worldwide average <sup>(4)</sup>           | 55  | 30-70  |  |   |
| RNPP site, Bangladesh (present study)      | 40  | 57.73  | 47.46 – 73.17  | Analysis of soil using gamma Spectrometry |

## CONCLUSION

The radioactivity and radiation levels in soil samples in and around the proposed Rooppur NPP Site must be useful as baseline data for the proposed NPP site at Rooppur for the purpose of detecting any changes in radioactivity in future. Radiological impact on the environment after commissioning of the nuclear power plant and any other nuclear fallout could be assessed by using this baseline data. The present study also helps to assess the radium equivalent activity, the absorbed dose rate, and external and internal radiation hazard due to presence of radionuclides in the soil. The report was submitted to the RNPP Implementation Committee.

## ACKNOWLEDGEMENTS

We express our sincere thanks and gratitude to the Chairman of Bangladesh Atomic Energy Commission. The authors wish to express their sincere thanks to Dr. Md. Shawkat Akbar, Project

Director of Rooppur Nuclear Power Plant for his sincere cooperation and providing all logistic support to carry out this work. The authors are thankful to employees of Health physics Division, AEC, Dhaka for their Cooperation.

**Conflicts of interest:** none to declare.

## REFERENCES

1. UNSCEAR, Sources, Effects and Risk of Ionization Radiation, United Nations Scientific Committee on the Effects of Atomic Radiation, (2000) Annex A: Exposure from Natural Sources, Report to General Assembly, UN, New York.
2. UNSCEAR, Sources, Effects and Risk of Ionization Radiation, United Nations Scientific Committee on the Effects of Atomic Radiation(1993) Report to General Assembly, UN, New York..
3. UNSCEAR, Sources, Effects and Risk of Ionization Radiation, United Nations Scientific Committee on the Effects of Atomic Radiation(1988) Report to General Assembly, UN, New York.
4. UNSCEAR, Sources, Effects and Risk of Ionization Radiation, United Nations Scientific Committee on the Effects of Atomic Radiation(1982) Report to General Assembly, UN, New York..
5. Mollah AS, Rahman MM, Hussain SR (1985) Preoperational background radiation level in and around 3 MW TRIGA Mark II research reactor, Report INST-19/HPRPD-3.

6. LEE MH and LEE CW (1997) Distribution and characteristics of  $^{239}$ ,  $^{240}$ Pu and  $^{137}$ Cs in the soil of Korea, *J. Environ Radioactivity*, **1**: 1–16.
7. Roy S, Hoque A, Begum M (2001) Distribution of  $^{137}$ Cs and Naturally Occurring Radionuclides in Soil at Sites of the Rooppur Nuclear Power plant. *Nuclear Science and Applications*. **10(1, 2)**: 33-38.
8. Beretka J and Mathew PJ (1985) Natural radioactivity of Australian building materials, industrial waste and by-products. *Health Physics*, **48(1)**: 87-95.
9. Delaune RD, Jones GL, Smith CJ (1986) Radionuclide Concentration in Louisiana Soils and Sediments. *Health Physics*, **51**: 239-244.
10. Noordin I (1999) Natural activities of  $^{238}$ U,  $^{232}$ Th and  $^{40}$ K in building materials. *Journal of Environmental Radioactivity*, **43**: 255-258.
11. Verga E, Bikit I, Slivka J et al., (2003) Danube sediment radioactivity investigation. *Proceedings of the Sixth International Symposium & Exhibition on Environment*.
12. Chowdhury MI, Alam MN, Hazari SKS (1999) Distribution of radionuclides in the river sediments and coastal soils of Chittagong, Bangladesh and evaluation of the radiation hazard. *Applied Radiation and Isotope*, **51**: 747-755.
13. Miah FK, Roy S, Touhiduzzaman M, Alam B (1998) Distribution of Radionuclides in soil samples in and around Dhaka city. *App Radiat Isot*, **49(1, 2)**: 133-137.
14. Chowdhury MI, Kamal M, Alam MN, Yeasmin Saleha, Mostafa MN (2006) Distribution of naturally occurring radionuclides in soils of the southern districts of Bangladesh. *Radiation Protection Dosimetry*, **118(1)**: 126-130.
15. Sharif AKM, Bilkis AB, Roy S, Sikder MDH, Idris Ali KM, Saifiullah S (1994) Concentrations of radionuclides in coastal sediments from the Bay of Bengal. *Sci Tot Environ*, **158**: 1-8.
16. Shenber MA (1997) Measurement of natural radioactivity levels in soil in Tripoli. *Appl Radiat Isot*, **48(1)**: 147-148.
17. Ajayi OS (2000) Distribution of natural radioactivity in rocks from Ikogoshi-Ekiti, southwestern Nigeria and its radiological implications. *Health Phy*, **79(2)**: 192-195.
18. Ali S, Tufail M, Jamie K, Ahmed A, Khan HA (1996) Gamma-ray activity and dose rate of brick samples from some area of North West frontier province (NWFP), Pakistan. *Science of Total Environment*, **187**: 247-252.
19. Uosif MAM and El-Taher A (2008) Radiological assessment of Abu-Tartur phosphate, western desert Egypt. doi: 10.1093/rpd/ncm502. *Radiation Protection Dosimetry*, **130**: 228-235.
20. Wang Z, He J, Du Y, He Y, Li Z, Chen Z, Chaowen Y (2012) Natural and artificial radionuclide measurements and radioactivity assessment of soil samples in eastern sichuan province (China). *Radiation Protection Dosimetry*, **150(3)**: 391–397.
21. Rahman MM, Rahman R, Begum M (2004) Distribution of natural and artificial radioactivity in soil from different districts of Bangladesh. *Journal of Bangladesh Academy of Sciences*, **28(1)**: 57-63.
22. Kabir KA, Islam SMA, Rahman MM (2009) Distribution of radionuclides in surface soil and bottom sediment in the district of Jessore, Bangladesh and evaluation of radiation hazard. *Journal of Bangladesh Academy of Science*, **1**: 117-130.
23. Sarkar SC, Ali I, Paul D, Bhuiyan MR, Islam SMA (2011) Measurement of natural and artificial radioactivity in soil at some selected Thanas around the TRIGA Mark-II research reactor at AERE, Savar, Dhaka. *Journal of Environmental Protection*, **2**: 1353-1359.