Identification of specific gene expression after exposure to low dose ionizing radiation revealed through integrative analysis of cDNA microarray data and the interactome

J.C. Son¹, H.O. Jeong¹, E.K. Lee^{1, 2}, S.G. No¹, D. Park^{3,4}, H.Y. Chung^{1,2*}

¹Interdisciplinary Research Program of Bioinformatics and Longevity Science, Pusan National University, Busan, Republic of Korea

²Molecular Inflammation Research Center for Aging Intervention (MRCA), College of Pharmacy, Pusan National University, Busan 46241, Republic of Korea

³Department of Predictive Toxicology, Korea Institute of Toxicology, Daejeon, Republic of Korea ⁴Center for Convergent Research of Emerging Virus Infection, Korea Research Institute of Chemical Technology, 141, Gajeong-ro, Yuseong-gu, Daejeon 34114, Republic of Korea

► Original article

*Corresponding authors: Hae Young Chung, PhD., Fax: +82 51 5188 2821 E-mail: hyjung@pusan.ac.kr

Revised: August 2017 **Accepted:** October 2017

Int. J. Radiat. Res., January 2019; 17(1): 15-23

DOI: 10.18869/acadpub.ijrr.17.1.15

ABSTRACT

Background: Accumulating reports suggest that the biological effects of low- and high- dose ionizing radiation (LDIR and HDIR) are qualitatively different and might cause different effects in human skin. Materials and Methods: To better understand the potential risks of LDIR, we analyzed three cDNA microarray datasets from the Gene Expression Omnibus database. Results: A pathway analysis showed that genes in immune-associated pathways were upregulated while those in cancer-associated pathways were downregulated in skin exposed to LDIR as compared with non-irradiated control skin. Consistently, according to a comparative gene ontology analysis, "antigen presentation and processing" was the most different gene ontology between the LIDR and HDIR transcriptomes. To identify key molecules regulated by LDIR, we constructed a protein-protein interaction network analysis using topological metrics. One of the key molecules with a high network scores was E1A binding protein p300 (EP300), which is a potential target of a new therapeutic strategy to promote anti-tumor immunity. Conclusion: Our results showed that LDIR exposure mainly induced the upregulation of immune-related genes including chemokines (CXCL1, CXCL2, and CXCL5) and interleukins (IL1B, IL11, IL6, IL15, and IL7). Additionally, LDIR induced the upregulation of antigen processing and presentation-related genes including CIITA, HLA-DQB1, and KIF26A, but these genes were downregulated in HDIR-exposed skin. Our protein network interaction results indicated that EP300 is downregulated by the immune response in skin after LDIR exposure.

Keywords: LDIR, cDNA microarray, transcriptome, Bioinformatics, EP300.

INTRODUCTION

According to the ICRP (International Commission on Radiological Protection), the incidence rate of cancer is proportional to the dose of radiation exposure, with no safety threshold (1, 2). In contrast, low dose ionizing

radiation (LDIR) effects, such as the adaptive response, the bystander effect, and genome instability are accepted, meaning that the LNT (the linear no threshold) model may not be correct for LDIR ^{(3-6).} The concept of an adaptive response refers to the idea that pre-exposure to LDIR can increase cellular tolerance to

subsequent radiation exposure through a hormetic mechanism and may thereby result in such consequences as life span prolongation ^(7,8), reduced tumor growth ⁽⁹⁾, improvement in degenerative disorders ⁽¹⁰⁻¹²⁾, and immune activation ⁽¹³⁾. Thus, LDIR-induced DNA damage below a certain threshold of severity could stimulate DNA repair systems, yielding a net long-term benefit to cellular functionality ⁽¹⁴⁾. Epigenetic modification ⁽¹⁵⁾ or differential gene expression ⁽¹⁶⁾ after LDIR exposure may contribute to this effect. Therefore, more studies need to be conducted to elucidate the effects of LDIR.

With the progression of high-throughput genomic technologies, global mRNA expression data is the most advanced current method for obtaining information about the cellular response to ionizing radiation. The use of gene ontology (GO) classifications of genes with statistically significant expression changes is a common procedure and a powerful tool for a deeper understanding of transcriptomics data. Functional genomic analysis contributes to the analysis of gene expression on a global scale. Therefore, we implemented a web application called Comparative GO to enable the comparison of the distribution of GO terms across multiple datasets ⁽¹⁷⁾. Although there were many previously existing web applications for GO analysis, none were suitable for the comparison of data from different sources. The collation of a tabular report containing a list of increased or decreased genes/proteins sorted with respect to their GO term contributes to understanding the effects of stimuli on biological pathways and the mechanisms of pathogenesis.

The purpose of this report was to assess the global cDNA expression profile of human skin samples exposed to LDIR and high dose ionizing radiation (HDIR) using cDNA microarray data sets. Using the result of microarray, we aimed to compare biological pathways and comparative GO terms between the skin samples exposed to LDIR and those exposed to HDIR. Skin is the first point of contact for radiation in most external exposure scenarios and skin samples can easily be obtained by biopsy to assess the biomolecular damage and biological response caused by LDIR.

significance of this study The is that immune-related and cancer-related biological pathways were meaningfully distinguishable between skin samples exposed to LDIR and those exposed to HDIR. Our integrative analysis of high-throughput gene expression data and subsequent construction of a protein-protein interaction (PPI) network provide evidence that better contributes to а understanding LDIR-induced immune enhancement.

MATERIALS AND METHODS

Microarray gene expression datasets

To identify differentially expressed genes through comparing skin samples (DEGs) exposed to LDIR with those exposed to HDIR, we screened publicly available microarray datasets and found the datasets GSE23807, GSE59861, and GSE29344 (total n = 25 subjects), which were retrieved from the Gene Expression Omnibus (GEO). In those three datasets, human skin samples (EpiDermFT-400, EPI-200, and AG01522 cells) were exposed to different total doses of radiation at dose rates between 10.5-50 cGy/min using a clinical X-ray irradiator. We defined LDIR as a total dose \leq 10 cGy and HDIR as a total dose \geq 100 cGy. The samples were harvested to obtain RNA 3-4 h after the radiation exposure.

Preprocessing for identification of genes with significant differences in expression

Before analyzing the microarray datasets, the gene expression values were log2 transformed, followed by normalization across all samples. Microarray gene expression values from samples representing different conditions (control vs. LDIR and control vs. HDIR) were compared by a two-class unpaired test using the Significance Analysis of Microarray tool (http:// statweb.stanford.edu/~tibs/SAM) to find DEGs. The DEGs in the LDIR and HDIR datasets were selected by an absolute fold-change of 1.5 (as compared with the control dataset) and a false discovery rate of 0.05. We then combined the DEGs obtained from the selected datasets.

16

Kyoto encyclopedia of genes and genomes (KEGG) pathway analysis

KEGG terms for biological pathways were as provided by the Database for used Visualization, Annotation, and Integrated Discovery (DAVID version 6.7) (https:// david.ncifcrf.gov/). Further. we conducted Fisher's exact *t*-test to detect the enrichment of DEGs within KEGG pathways (p < 0.05).

Comparative GO analysis

To compare GO terms between the LDIR and HDIR transcriptomes, we conducted an analysis Comparative GO using (http:// www.comparativego.com). Other GO web applications can compare a single sample to another reference sample and present differentially regulated GOs with *p*-values. However, Comparative GO can compare the distribution of GO terms among gene expression datasets from several different sources and provides a tabular report of the number of genes exceeding the fold-change threshold among treatment groups, enabling the identification of the GO terms that contain the most DEGs.

Analysis of PPI Network

To identify proteins that acted as hubs in the PPI network, we used the HPRD (Human Protein Reference Database) release 9 and IntAct molecular interaction database) version 2.0. (http://www.cytoscape.org/) was Cytoscape used for visualizing the interaction network. The degree and betweenness centrality were calculated for each node in the PPI network (18). The degree is classified as the number of interacting partners that connect each protein to its neighbors, while betweenness centrality represents the number of short paths between one protein and other proteins. These two topological parameters have been known to assist the detection of functionally important genes. Clustering analysis was also conducted to distinguish groups that have high degrees and betweenness centralities compared with the other groups.

RESULTS

Identification of DEGs Associated with LDIR and HDIR Exposure Levels

To identify radiation-regulated DEGs using three human skin datasets from investigations in which different radiation doses were administered, we selected three **cDNA** microarrav datasets representing LDIR exposure, HDIR exposure, and non-irradiated control skin. As a result, we obtained 1,833 DEGs, which we collectively termed the LDIR and HDIR transcriptomes. There were 844 DEGs in the LDIR dataset and 989 DEGs in the HDIR dataset relative to the non-irradiated control (figure 1). In the LDIR transcriptome, 311 genes were increased and 533 were decreased. In the HDIR transcriptome, 387 genes were increased and 602 were decreased. A total of 352 DEGs were shared between the LDIR and HDIR transcriptomes.

Sample clustering analysis

In order to compare the gene expression patterns among samples, we utilized MeV software ⁽¹⁹⁾. As illustrated in figure 2, we used hierarchical link clustering (HLC) and principal component analysis (PCA) methods. Based on HLC analysis, the samples were divided into two clusters: control and radiation. Furthermore, the radiation samples were divided into two clusters: LDIR and HDIR (figures 2a & b). PCA was employed to determine the gene specificity among different groups. Irradiated samples were clearly separated from the control group and LDIR samples showed distinct expression patterns from those of HDIR samples (figure 2c). These results suggested that high-throughput gene expression profiling can differentiate between LDIR and HDIR exposure.

Pathway Enrichment Analysis of the LDIR and HDIR Transcriptomes

We performed a KEGG pathway enrichment analysis using the DAVID database, which showed that the upregulated genes in the LDIR

transcriptome mainly related were to immune-associated pathways such as "cvtokine-cvtokine receptor interaction". "apoptosis", and "NOD-like receptor signaling pathway". In addition, LDIR exposure caused the down-regulation of pathways related to the process of skin cancer, such as "melanogenesis", "basal cell carcinoma", "hedgehog signaling pathway", and "pathways in cancer" (table 1).

In the HDIR transcriptome, the pathways associated with upregulated genes were largely cancer-related. including "p53 signaling pathway", "ErbB signaling pathway", "VEGF signaling pathway". "focal adhesion", "ECM-receptor interaction", and several cancer pathways. In addition, downregulated genes related to cell proliferation presented a bias toward pathways such as "cell cycle", "meiosis", and "DNA replication" (table 2).

Comparative GO analysis of LDIR and HDIR transcriptomes

We applied the Comparative GO web application to compare the lists of GO terms between the LDIR and HDIR transcriptomes. As shown in Figure 3, immune-related biological processes, such as antigen processing and presentation (101.5 times), immune response (24.5 times), and regulation of tumor necrosis factor super family cytokine production (8.6 times) showed differences between the upregulated genes in LDIR and HDIR transcriptomes. The specific genes responsible for antigen processing and presentation that were upregulated by LDIR exposure but downregulated by HDIR transcriptomes were major histocompatibility complex class II, DQ beta 1 (HLA-DOB1), class Π major histocompatibility complex transactivator (*CIITA*), and kinesin family member 6° A (*KIF26A*).

Target molecules in the PPI network

To determine key molecules that regulate the immune response to LDIR exposure, we conducted a PPI network analysis because the PPI network has been widely used to develop drug targets, disease genes, and essential genes in organisms ranging from yeast to humans. In this study, information on human PPI networks was obtained from the HPRD and IntAct databases. As a result, the PPI network of the significantly expressed genes in the LDIR transcriptome was composed of 184 genes (77 upregulated genes and 107 downregulated genes), as shown in Figure 4a. To identify target molecules in the PPI network, we then performed a clustering analysis using the degree and betweenness centrality as the topological parameters. As a result, E1A binding protein p300 (EP966) was predicted as a target molecule because it had a higher degree and betweenness centrality than the other nodes (figure 4b).



Figure 1. Differentially expressed genes (DEGs) in the low-dose ionizing radiation (LDIR)- and high-dose ionizing radiation (HDIR)-exposed skin transcriptomes. The DEGs were selected using the absolute cut-off criteria of a fold-change > 1.5 and a false discovery rate < 0.05. As a result, we obtained 1,833 DEGs from the LDIR and HDIR transcriptomes (844 in LDIR and 989 in HDIR).

Int. J. Radiat. Res., Vol. 17 No. 1, January 2019



Figure 2. Visualizations of cDNA microarray gene expression profiles of the LDIR- and HDIR-exposed skin transcriptomes.
(a) Analysis of microarray data shown as a heat map comparing global gene expression patterns among the control, LDIR, and HDIR transcriptomes. Red and green colors indicate differentially up- or down-regulated genes, respectively (fold-change > 2).
(b) Hierarchical link clustering and (c) principal component analysis plot. Two-dimensional scatter plots for dividing the samples based on two principal components are shown in (c).

Category	Term	Count	p -value
Upregulated			
KEGG_PATHWAY	Cytokine-cytokine receptor interaction	17	1.53E-04
KEGG_PATHWAY	Apoptosis	8	2.7E-03
KEGG_PATHWAY	NOD-like receptor signaling pathway	6	1.1E-02
Downregulated			
KEGG_PATHWAY	Complement and coagulation cascades	7	5.4E-03
KEGG_PATHWAY	Melanogenesis	8	8.5E-03
KEGG_PATHWAY	Nitrogen metabolism	4	1.6E-02
KEGG_PATHWAY	Basal cell carcinoma	5	3.9E-02
KEGG_PATHWAY	Hedgehog signaling pathway	5	4.2E-02
KEGG_PATHWAY	Pathways in cancer	14	4.4E-02

Table 2. KEGG pathway	/ analysis of up- or	down-regulated D	EGs in HDIR.
-----------------------	----------------------	------------------	--------------

Category	Term	Count	<i>p</i> -value
Upregulated			
KEGG_PATHWAY	p53 signaling pathway	8	5.3E-04
KEGG_PATHWAY	ErbB signaling pathway	8	2.3E-03
KEGG_PATHWAY	VEGF signaling pathway	6	2.1E-02
KEGG_PATHWAY	Focal adhesion	10	2.6E-02
KEGG_PATHWAY	Non-small cell lung cancer	5	2.7E-02
KEGG_PATHWAY	Small cell lung cancer	6	3.2E-02
KEGG_PATHWAY	ECM-receptor interaction	6	3.2E-02
KEGG_PATHWAY	Pathways in cancer	13	4.4E-02
Downregulated			
KEGG_PATHWAY	Cell cycle	20	7.58E-07
KEGG_PATHWAY	Oocyte meiosis	13	2.0E-03
KEGG_PATHWAY	DNA replication	6	1.6E-02

Int. J. Radiat. Res., Vol. 17 No. 1, January 2019



Figure 3. Comparison between the gene ontology (GO) terms associated with upregulated genes in the LDIR and HDIR transcriptomes. Protein enrichment values were normalized based on the number of genes in each sample. The enrichment values of the smaller samples are scaled higher to be comparable to those of the bigger samples. Among the DEGs in the LDIR and HDIR transcriptomes, those whose protein enrichment value underwent a rate of change > 2 were selected and categorized according to their biological process. The rate of change reflects the geometric average of the enrichment value fold-change among samples in that GO group.



Figure 4. Protein-protein interaction network composed of DEGs in the LDIR transcriptome. (a) The network, in which the nodes indicate proteins and the edges indicate the interactions between the proteins inferred from the IntAct and HPRD databases. A total of 77 proteins that were upregulated in the LDIR transcriptome are represented in red, and 107 proteins that were downregulated in the LDIR transcriptome is correlated with the number of interacting partners and BC (betweenness centrality) represents the number of short paths between one protein and other proteins.

20

DISCUSSION

Even though the molecular mechanisms underlying the LDIR-induced immune response are still unknown, low dose radiotherapy is considered as an alternative option for the treatment of inflammatory diseases and degenerative joint disorders in Germany and other European countries ⁽²⁰⁻²³⁾. Interestingly, LDIR has been reported to decrease the incidence rates of several cancers, including B16 melanoma ⁽²⁴⁾.

In accordance with these findings, our results showed that among the DEGs identified in the LDIR transcriptome, most of them were upregulated immune response genes including chemokines (CXCL7, CXCL2, and CXCL5) and interleukins (IL7B, IL11, IL6, IL15, and IL7). Chemokines have important functions in wound healing. lymphocyte homing, chemotactic activity for lymphocytes, and influencing the overall type 1/type 2 balance of an immune response ^(25, 26). Akira *et al.* found that after a single low dose exposure (1.0 cGy), chemokines (CXCL1, CXCL2, and CXCL6) were significantly upregulated in normal human fibroblasts (27). Interleukins are mediator of inflammation, the immune system, and cancer. The increases of IL1B and IL6 caused by 4.5 Gy can contribute to the enhancement of immune responses (28). Furthermore, an increased expression level of IL15 was reported following irradiation; this interleukin acts through the selective inhibition of tumor-promoting molecules, and has been considered as a potential new compound for use in cancer treatment ⁽²⁹⁾. The T cell lymphocyte development-related cytokine IL3 was upregulated in low dose exposed mice ⁽³⁰⁾. These data imply that LDIR-induced immune responses are likely to be activated to eliminate damaged cells as part of a tumor inhibition, and this might lead to decreases in several cancer-related pathways as indicated by our KEGG analysis (table 1).

Comparative GO analysis revealed that LDIR induced the up-regulation of antigen processing- and presentation-related genes including *CIITA*, *HLA-DQB1*, and *KIF26A*, but HDIR was associated with the down-regulation

Int. J. Radiat. Res., Vol. 17 No. 1, January 2019

of these genes. Efficient antigen presentation and processing is important for inducing potent anti-tumor immune responses because the function of the MHC class molecules is critical in some cancers ⁽³¹⁻³⁴⁾. Kinesin is a motor protein that is related to the transport of MHC class II containing complexes along microtubules to the late endosomal compartment. Recently, many reports have revealed that aberrant gene expression of kinesins plays a key role in a variety of human cancers, suggesting that kinesins may represent new molecular targets for cancer therapy (35-37). MHC class II is important for the activation of helper T cells. It is well documented that helper T cells contribute to anti-tumor immunity (38). Several solid cancers do not express MHC class II, and the involvement of helper T cells depends on infiltrating antigen-presenting cells. CIITA, the master regulator of MHC class II transcription, is a non-DNA binding protein and has also been found to be increased during immune responses. Therefore, we propose that *KIF82A*, *HLA-DOB1*, and CIITA play important roles in the LDIR induced effects on antigen presentation and processing.

An analysis of the hub genes among the DEGs in the LDIR dataset (figure 4a) revealed that EP300 had the highest connective degree. EP300 is involved in several processes including proliferation, apoptosis, and DNA damage (39, 40). Yujie *et al.* reported that *EP966* plays an important role in anti-tumor immunity (41), and inhibitors of EP300 were well-established caloric restriction mimetics ⁽⁴²⁾. Acetyl coenzyme А induces autophagy through а transcription-independent process related to the reduction of acetyltransferase EP300 (43).

In summary, our findings show differences in signaling pathways and biological processes between the transcriptomes of human skin samples exposed to LDIR and those exposed to HDIR. Our findings are as follows: (1) the molecular response after LDIR involves the up-regulation of the immune response by the secretion of immune-related cytokines; (2) "antigen presentation and processing" is the GO term with the greatest difference between the LDIR and HDIR transcriptomes; and (3) the

down-regulation of EP300 caused by LDIR could lead to an inhibition of cancer development through anti-tumor immunity. These results provide a better understanding of the risks of using LDIR in the clinical setting and specifically highlight its effects on the human immune system.

ACKNOWLEDGEMENT

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (2009-0083538). In addition, this work was supported by the National Research Foundation of Korea Grant funded by the Korea Government (NRF-2015M3A9B8029074). The authors are grateful to the Aging Tissue Bank (Busan, Korea) for supplying the research informations used in this work.

Conflicts of interest: Declared none.

REFERENCES

- Huang L, Snyder AR, Morgan WF (2003) Radiation-induced genomic instability and its implications for radiation carcinogenesis. *Oncogene*, 22(37): 5848-5854.
- Little J (1998) Radiation-induced genomic instability. Int J Radiat Biol, 74(6): 663-671.
- 3. Wall B, Kendall G, Edwards A, Bouffler S, Muirhead C, Meara J (2014) What are the risks from medical X-rays and other low dose radiation? *The British Journal of Radiology*.
- Hoffmann G and Stempsey W (2008) The hormesis concept and risk assessment: are there unique ethical and policy considerations? *Human & Experimental Toxicology*, 27(8): 613-620.
- Matsumoto H, Tomita M, Otsuka K, Hatashita M (2009) A new paradigm in radioadaptive response developing from microbeam research. *Journal of Radiation Research*, 50 (Suppl A): A67-A79.
- Lorimore SA, Coates PJ, Wright EG (2003) Radiationinduced genomic instability and bystander effects: interrelated nontargeted effects of exposure to ionizing radiation. Oncogene, 22(45):7058-7069.
- Moskalev A, Plyusnina E, Shaposhnikov M (2011) Radiation hormesis and radioadaptive response in Drosophila melanogaster flies with different genetic backgrounds: the role

of cellular stress-resistance mechanisms. *Biogerontology*, **12(3)**: 253-263.

- Seong KM, Kim CS, Seo S-W, Jeon HY, Lee B-S, Nam SY, Yang KH, Kim J-Y, Kim CS, Min K-J (2011) Genome-wide analysis of low-dose irradiated male Drosophila melanogaster with extended longevity. *Biogerontology*, 12(2): 93-107.
- Shin SC, Lee KM, Kang YM, Kim K, Lim SA, Yang KH, Kim JY, Nam SY, Kim HS (2011) Differential expression of immuneassociated cancer regulatory genes in low-versus highdose-rate irradiated AKR/J mice. *Genomics*, *97(6):* 358-363.
- Micke O, Seegenschmiedt MH, Diseases GCGoRfB (2004) Radiotherapy in painful heel spurs (plantar fasciitis) results of a national patterns of care study. *Int J Radiat Oncol Biol Phys*, 58(3): 828-843.
- 11. Niewald M, Fleckenstein J, Naumann S, Ruebe C (2007) Long-term results of radiotherapy for periarthritis of the shoulder: a retrospective evaluation. *Radiat Oncol*, **2**: 34.
- Adamietz B, Schulz-Wendtland R, Alibek S, Uder M, Sauer R, Ott O, Keilholz L (2010) Calcifying Tendonitis of the Shoulder Joint. *Strahlentherapie und Onkologie*, **186(1)**: 18 -23.
- Ina Y and Sakai K (2005) Activation of immunological network by chronic low-dose-rate irradiation in wild-type mouse strains: analysis of immune cell populations and surface molecules. Int J Radiat Biol, 81(10): 721-729.
- 14. Rödel F, Kamprad F, Sauer R, Hildebrandt G (2002) Functional and molecular aspects of anti-inflammatory effects of low-dose radiotherapy. *Strahlentherapie und Onkologie*, **178(1)**: 1-9.
- Ma S, Liu X, Jiao B, Yang Y, Liu X (2010) Low-dose radiation -induced responses: focusing on epigenetic regulation. Int J Radiat Biol, 86(7): 517-528.
- Rodel F, Frey B, Gaipl U, Keilholz L, Fournier C, Manda K, Schollnberger H, Hildebrandt G, Rodel C (2012) Modulation of inflammatory immune reactions by low-dose ionizing radiation: molecular mechanisms and clinical application. *Current Medicinal Chemistry*, **19(12)**: 1741-1750.
- Fruzangohar M, Ebrahimie E, Ogunniyi AD, Mahdi LK, Paton JC, Adelson DL (2013) Comparative GO: a web application for comparative gene ontology and gene ontologybased gene selection in bacteria. *PloS one*, **8(3)**: e58759.
- Newman ME (2005) A measure of betweenness centrality based on random walks. Social Networks, 27(1): 39-54.
- Saeed A, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M (2003) TM4: a free, open-source system for microarray data management and analysis. *Biotechniques*, **34(2)**:374.
- Hildebrandt G, Radlingmayr A, Rosenthal S, Rothe R, Jahns J, Hindemith M, Rödel F, Kamprad F (2003) Low-dose radiotherapy (LD-RT) and the modulation of iNOS expression in adjuvant-induced arthritis in rats. *Int J Radiat Biol*, **79** (12): 993-1001.
- Seegenschmiedt MH, Micke O, Willich N (2004) Radiation therapy for nonmalignant diseases in Germany. Strahlentherapie und Onkologie, 180(11): 718-730.

Int. J. Radiat. Res., Vol. 17 No. 1, January 2019

- Heyd R, Tselis N, Ackermann H, Röddiger SJ, Zamboglou N (2007) Radiation therapy for painful heel spurs. *Strahlentherapie und Onkologie*, **183(1)**: 3-9.
- Hildebrandt JJ, Hindemith M, Spranger S, Sack U, Kinne RW, Madaj-Sterba P, Wolf U, Kamprad FG (2000) Effects of low dose radiation therapy on adjuvant induced arthritis in rats. *Int J Radiat Biol*, **76(8)**: 1143-1153.
- 24. Lugade AA, Moran JP, Gerber SA, Rose RC, Frelinger JG, Lord EM (2005) Local radiation therapy of B16 melanoma tumors increases the generation of tumor antigen-specific effector cells that traffic to the tumor. *The Journal of Immunology*, **174(12)**: 7516-7523.
- 25. D'Ambrosio D, Panina-Bordignon P, Sinigaglia F (2003) Chemokine receptors in inflammation: an overview. *Journal of Immunological Methods*, **273(1)**: 3-13.
- Rossi D and Zlotnik A (2000) The biology of chemokines and their receptors. Annual Review of Immunology, 18(1): 217-242.
- Fujimori A, Okayasu R, Ishihara H, Yoshida S, Eguchi-Kasai K, Nojima K, Ebisawa S, Takahashi S (2005) Extremely low dose ionizing radiation up-regulates CXC chemokines in normal human fibroblasts. *Cancer Research*, 65(22): 10159 -10163.
- Hosoi Y, Miyachi H, Matsumoto Y, Enomoto A, Nakagawa K, Suzuki N, Ono T (2001) Induction of interleukin-1β and interleukin-6 mRNA by low doses of ionizing radiation in macrophages. *International Journal of Cancer*, *96(5):* 270-276.
- Shanmugham L, Petrarca C, Frydas S, Donelan J, Castellani M, Boucher W, Madhappan B, Tete S, Falasca K, Conti P (2006) IL-15 an immunoregulatory and anti-cancer cytokine. Recent advances. *Journal of Experimental & Clinical Cancer Research: CR*, 25(4): 529-536.
- 30. Van Roon J, Glaudemans K, Bijlsma J, Lafeber F (2003) Interleukin 7 stimulates tumour necrosis factor α and Th1 cytokine production in joints of patients with rheumatoid arthritis. Annals of the Rheumatic Diseases, **62(2)**: 113-119.
- Steidl C, Shah SP, Woolcock BW, Rui L, Kawahara M, Farinha P, Johnson NA, Zhao Y, Telenius A, Neriah SB (2011) MHC class II transactivator CIITA is a recurrent gene

fusion partner in lymphoid cancers. *Nature*, **471(7338)**: 377-381.

- Poschke I, Mougiakakos D, Kiessling R (2011) Camouflage and sabotage: tumor escape from the immune system. *Cancer Immunology, Immunotherapy*, 60(8): 1161-1171.
- Corthay A, Skovseth DK, Lundin KU, Røsjø E, Omholt H, Hofgaard PO, Haraldsen G, Bogen B (2005) Primary antitumor immune response mediated by CD4+ T cells. *Immunity*, 22(3): 371-383.
- Nakanishi Y, Lu B, Gerard C, Iwasaki A (2009) CD8+ T lymphocyte mobilization to virus-infected tissue requires CD4+ T-cell help. *Nature*, 462(7272): 510-513.
- 35. Yu Y and Feng YM (2010) The role of kinesin family proteins in tumorigenesis and progression. *Cancer*, **116(22)**: 5150-5160.
- Rath O and Kozielski F (2012) Kinesins and cancer. Nature Reviews Cancer, 12(8): 527-539.
- Huszar D, Theoclitou ME, Skolnik J, Herbst R (2009) Kinesin motor proteins as targets for cancer therapy. *Cancer and Metastasis Reviews*, 28(1-2): 197-208.
- Quezada SA and Peggs KS (2011) Tumor-reactive CD4+ T cells: plasticity beyond helper and regulatory activities. *Immunotherapy*, 3(8): 915-917.
- Chan HM and La Thangue NB (2001) p300/CBP proteins: HATs for transcriptional bridges and scaffolds. *Journal of Cell Science*, **114(13)**: 2363-2373.
- Goodman RH and Smolik S (2000) CBP/p300 in cell growth, transformation, and development. *Genes & Development*, 14(13): 1553-1577.
- Liu Y, Wang L, Predina J, Han R, Beier UH, Wang L-CS, Kapoor V, Bhatti TR, Akimova T, Singhal S (2013) Inhibition of p300 impairs Foxp3+ T regulatory cell function and promotes antitumor immunity. *Nature Medicine*, **19(9)**: 1173-1177.
- Pietrocola F, Pol J, Vacchelli E, Rao S, Enot DP, Baracco EE, Levesque S, Castoldi F, Jacquelot N, Yamazaki T (2016) Caloric restriction mimetics enhance anticancer immunosurveillance. *Cancer Cell*, **30(1)**: 147-160.
- Mariño G, Pietrocola F, Eisenberg T, Kong Y, Malik SA, Andryushkova A, Schroeder S, Pendl T, Harger A, Niso-Santano M (2014) Regulation of autophagy by cytosolic acetyl-coenzyme A. *Molecular Cell*, 53(5): 710-725.

[Downloaded from ijrr.com on 2025-08-23]