

Measurement of radium-226 concentration and dose calculation of drinking water samples in Guilan province of Iran

A. Abbasi^{1,2*} and V. Bashiry³

¹Nuclear Science and Technology Institute, (NSTI), Tehran, Iran

²Faculty of Engineering, University of Kyrenia, Kyrenia, North Cyprus, Mersin 10, Turkey

³Faculty of Engineering, Cyprus International University, Nicosia, Northern Cyprus, Mersin 10, Turkey

ABSTRACT

Background: Radium-226 (²²⁶Ra) is a product of the ²³⁸U radionuclide decay series that significantly incorporated into the human body through water intake. It can also potentially cause a series of health problems including cancers of the digestive system. Radium-226 (²²⁶Ra) is a product of the ²³⁸U radionuclide decay series that significantly incorporated into the human body through water intake. It can also potentially cause a series of health problems including cancers of the digestive system. **Materials and Methods:** ²²⁶Ra has been determined in drinking water samples collected from various locations of Guilan province of Iran. The water samples are taken from public water and urban tap water sources. For evolution of gross α and β exposure precipitation method and proportional scintillator system was used. The radon emanation method was used to measure the radium concentration in drinking water. **Results:** The measurements showed the gross α and β concentration ranges were between $<38 \text{ mBq l}^{-1}$ to 92 mBq l^{-1} and $<41 \text{ mBq l}^{-1}$ to 328 mBq l^{-1} , respectively. The radium concentration range was between 2 mBq l^{-1} to 38.2 mBq l^{-1} . The resulting contribution to the annual effective dose due to the digestion of ²²⁶Ra in water was calculated to be between $<0.4 \mu\text{Sv y}^{-1}$ to $7.8 \mu\text{Sv y}^{-1}$, respectively. **Conclusion:** The average concentration of ²²⁶Ra was found (7.6 mBq l^{-1}) in drinking water samples and the average annual effective dose, from the digestion of ²²⁶Ra in water samples was calculated to be $1.5 \mu\text{Sv y}^{-1}$. According to UNSCEAR, the annual effective dose value by ingestion is report to be 0.12 mSv y^{-1} .

Keywords: Radium-226, ingestion, drinking water, precipitation method.

► Short report

*Corresponding author:

Dr. Akbar Abbasi,

Fax: +90 392 815 99 36

E-mail: akbar.abbasi@aeoi.org.ir
akbar.abbasi@Kyrenia.edu.tr

Revised: Dec. 2015

Accepted: Jan. 2016

Int. J. Radiat. Res., October 2016;
14(4): 361-366

DOI: 10.18869/acadpub.ijrr.14.4.361

INTRODUCTION

The main sources of natural radioactivity are ²³⁸U, ²³²Th and their decay products and ⁴⁰K. Radium-226 is a product of the ²³⁸U radionuclide decay series with half-life 1600 y that emits alpha and beta particles. Among radionuclides present in groundwater, the most radiotoxic and of concern to human health is radium (¹). The ²²⁶Ra concentration in surface waters is commonly scant, but groundwater sources can contain a considerable concentration dependent

upon the uranium and thorium content of the surrounding geology. Radium can enter groundwater by dissolution of aquifer materials that desorption from rock or sediment surfaces and ejection from minerals by radioactive decay.

Long term exposure to elevated levels of radium in drinking water has been associated with an increased risk of bone cancer development. Although the consumption of drinking water containing radium concentrations below the established MCL (Minimum Concentration Limit $\approx 1 \text{ Bq l}^{-1}$) does

not alleviate all potential risk. Altogether the impact of public health at these concentrations is considered negligible (2).

Since the study of radioactivity concentration in drinking water has not already been done in this area and also because of the high percentage of various diseases, including cancer of the digestive system in the studied area, this study is performed to obtain an estimate of the activity concentration of ²²⁶Ra in public water and urban tap water sources. Furthermore, the radiation dose from ingestion of drinking water by public in Guilan province of Iran was estimated.

MATERIALS AND METHODS

In this study, 28 drinking water samples from the waters which resource public waters (wells) and spring waters were collected. The study area was north section of Guilan province (figure 1). For sample preparation, the Environmental Protection Agency (EPA) protocol (3) was adopted, in which samples were collected in 4 Li container containing a solution with pH≈2. First of all, each sample was measured the gross a and b exposure by using a gas scintillation proportional counter (CANBERRA) to determine the ²²⁸Ra level according EPA protocol. Commercial software Genie 2000 v2.1 was used for data analysis. The ²²⁶Ra in the drinking water sample was concentrated and separated in solution by precipitation method. The

precipitate obtained was dissolved in EDTA reagent, placed in a sealed bubbler and stored for ingrowth of ²²²Rn. After ingrowth ²²²Ra, the gas was purged into a scintillation Lucas cells, by noble gas. When the ²²²Rn daughters are in equilibrium with the parent (approximately, 4h), the scintillation Lucas cell, was counted by using a Pylon AB-5 radon field-portable scintillation measurement system. Use of the Lucas cell for radon measurement is an established approach in field and laboratory settings (4-7). The separation procedure and instrumentals' calibration was performed using a standard solution of this nuclide. The Lucas cell was evacuated and then attached to a sample cylinder with in-line valves, expansion tubing, and a pressure gage. The Lucas cell was filled with sample by opening the cylinder to the evacuated expansion tubing and then closing the cylinder valve and opening the valve to the Lucas cell. After the pressure stabilized, the process was repeated if necessary until the cell was filled with natural gas to approximately atmospheric pressure. The final pressure was recorded and the Lucas cell placed on the AB-5 photomultiplier tube to count the total alpha activity. The software SP-55 version 1.3 was used for data analysis. The counting interval was 90 minutes and was repeated as time p ermitted. Radon activities were calculated from the counting data after correcting for the efficiency of the Lucas cell, the cell volume, the recorded pressures, and the time interval between sample




Figure 1. The study area in Guilan province of Iran.

collections and counting.

Radium estimation in drinking water

The ²²⁶Ra activity concentration was calculated in the pCi per liter using equation (1):

$$C_{Ra} = \frac{C \lambda t_3}{2.22 EV(1 - e^{-\lambda t_1})(e^{-\lambda t_2})(1 - e^{-\lambda t_3})} \quad (1)$$

Where; C_{Ra} is the ²²⁶Ra activity concentration (pCi l⁻¹), and C, E and V are the net count rate (cpm), calibration constant and sample volume (Li), respectively. The t₁, t₂, t₃ and λ parameters are the passed time between the first and second de-emanations (min), the time between the second de-emanation and counting, the counting time and the decay constant of ²²²Rn (1.26 × 10⁻⁴ min⁻¹), respectively. The conversion factor constant from dpm/pCi is 2.22 (8).

The calibration constant (E), was calculated by using equation (2) as follow:

$$E = \frac{C}{A(1 - e^{-\lambda t_1})(e^{-\lambda t_2})} \quad (2)$$

where; C_is the net count rate (cpm), A is the activity of ²²⁶Ra in the bubbler (dpm), t₁ and t₂ are ingrowth time of ²²²Rn and the decay time of ²²²Rn occurring between de-emanation and counting (min), respectively (3).

Radium dose calculation

The dose calculation from radium as a result of drinking water is calculated by equation (3) as follow:

$$D = D_{WI} \times D_{CF} \times Y \times C_R \quad (3)$$

where; D is the annual effective dose (Sv y⁻¹) from the digestion of ²²⁶Ra in water, C_R is the radium concentration in water (Bq l⁻¹), D_{WI} is the daily consumed of water (estimated as 2l d⁻¹), D_{CF} is the dose conversion factor for adults, (2.8 × 10⁻⁷Sv Bq⁻¹) and Y is year (365 d) (9). In all calculations the uncertainties are given within 1 standard deviation.

RESULTS

A total of 28 samples that were taken from the study area were included 14 tap water samples, 12 well water samples and 2 spring

water samples. The results of gross a and b concentration in the drinking water samples are shown in table 1. The gross a and b range were between the low limit of detection LLD (<38 mBq l⁻¹) to 92 mBq l⁻¹ and LLD (<41 mBq l⁻¹) to 328 mBq l⁻¹, respectively. Also the ²²⁶Ra concentration and annual effective dose contribution due to the digestion of water samples used in North Guilan Iran are shown in table 1. The ²²⁶Ra concentrations for 8 samples were below the LLD=2 mBq l⁻¹ and maximum concentration was 38.2 mBql⁻¹ in WS-8 (Khotbeh Sera). The annual effective dose range was calculated between to be <0.4 μSv y⁻¹ to 7.8 μSv y⁻¹. The highest values related to tap water sample WS-8 (Khotbeh Sera) and spring water sample HSWS-4 (Kote Kumeh).

The correlation parameter between the radium-226 concentration and water pH are shown in figure 2. The result showed that high ²²⁶Ra concentrations were in alkaline water samples.

In following the cause of high grossa, gross b and radium concentration area, with the geological map, the veins mass of metamorphic minerals and pyroxene andesitic contain were found.

DISCUSSION

The screening levels for gross α and gross β in drinking water samples were below 500 mBq l⁻¹ and 1000 mBq l⁻¹, respectively, therefore, below the recommended World Health Organization (WHO) guideline level for drinking water (1.0 Bq l⁻¹) (10). Also, according to the Environmental Protection Agency (EPA) the ²²⁶Ra concentration in drinking water as safe limit is at the level of 0.74 Bql⁻¹ (3). This implies that all the water samples had values below the recommended levels.

In this study the average annual effective dose in water samples area was less than the WHO guideline level value of 0.1 mSy y⁻¹(10). Also the ²²⁶Ra concentrations measured in this study were compared with other published papers. These results were within reported ranges as shown in table 2.

Table 1. The gross a, gross b, radium-226 concentration and annual effective dose from the digestion of ²²⁶Ra in water samples used in North Guilan province, Iran.

No	Sample Code	Regain Name	Sample Resource	pH	Gross Alpha (mBq l ⁻¹)	Gross Beta (mBq l ⁻¹)	²²⁶ Ra Concentration (mBq l ⁻¹)	Annual dose due to Ingestion of ²²⁶ Ra (μSv y ⁻¹)
1	WS-1	Astara	Tap Water	7.6	>LLD*	>LLD**	>LLD***	0.4>
2	WS-2	Abas Abad	Tap Water	7.4	>LLD	4 ±54	0.03±7.4	1.5
3	WS-3	Lavandvil	Tap Water	7.1	>LLD	>LLD	0.02±2.1	0.4
4	HSWS-4	KoteKumeh	Spring Water	8.8	5±81	8±310	0.08±24.8	5.1
5	WS-5	Chooabar	Well Water	7.2	>LLD	2±61	0.06±3.4	0.7
6	WS-6	Vizneh	Tap Water	6.9	>LLD	3± 42	0.02±2.7	0.6
7	WS-7	Haviq	Tap Water	7.2	>LLD	4±71	0.04±4.6	0.9
8	WS-8	Khotbeh Sera	Tap Water	7.6	6±92	12±328	0.08±38.2	7.8
9	WS-9	Lisar	Tap Water	7.2	>LLD	>LLD	>LLD	0.4>
10	WS-10	Jemakoooh	Well Water	6.9	4±68	>LLD	0.07±8.2	1.7
11	WS-11	Talesh	Tap Water	7.1	>LLD	2±49	0.02±11.4	2.3
12	WS-12	Asalem	Tap Water	7.2	>LLD	>LLD	>LLD	0.4>
13	WS-13	Gisum	Well Water	7.4	>LLD	>LLD	>LLD	0.4>
14	HSWS-14	Laktoshan	Spring Water	7.7	3±78	4±75	0.09±21.5	4.4
15	WS-15	Paresar	Well Water	7.2	>LLD	>LLD	0.02±3.4	0.7
16	WS-16	Ardjan	Well Water	7.5	2±44	6±97	0.04±14.3	2.9
17	WS-17	Rezvanshahr	Tap Water	7.2	>LLD	>LLD	>LLD	0.4>
18	WS-18	Punel	Tap Water	7.5	>LLD	>LLD	>LLD	0.4>
19	WS-19	Shanderman	Well Water	7.5	>LLD	14±257	0.02±10.8	2.2
20	WS-20	Masal	Well Water	7.3	2±43	12±263	0.05±2.4	0.5
21	WS-21	Ziaber	Well Water	6.9	>LLD	4±88	0.03±5.2	1.1
22	WS-22	Bandar Anzali	Tap Water	7.2	>LLD	>LLD	0.04±5.3	1.1
23	WS-23	Abkenar	Well Water	7.6	>LLD	7±145	0.04±3.5	0.7
24	WS-24	Sowme'eh Sara	Well Water	7.5	3±51	4±57	0.03±21.3	4.4
25	WS-25	Fuman	Tap Water	7.4	>LLD	>LLD	>LLD	0.4>
26	WS-26	Khomam	Well Water	7.6	>LLD	3±48	>LLD	0.4>
27	WS-27	Rasht (Pirbazar)	Well Water	7.4	>LLD	12±281	0.05±2.8	0.6
28	WS-28	Rasht	Tap Water	7.2	>LLD	5±81	0.02±3.3	0.7
	Range			-6.9 8.8	>LLD- 92	>LLD-328	>LLD - 38.2	7.8 - 0.4>

*LLD= 38 mBq l⁻¹; **LLD=41 mBq l⁻¹; *** LLD=2 mBq l⁻¹




Figure 2. The correlation parameter between the radium-226 concentration and water pH.

Table 2. The radium-226 concentration in drinking water in the world.

Location	Sample resource	²²⁶ Ra concentration	Reference
Shandiz, Iran	Springs and rivers	MDA–2.2 Bq l ⁻¹	Binesh <i>et al.</i> ⁽¹¹⁾
Extramadura, Spin	Springs, wells and spas	<0.2–142 mBq l ⁻¹	Lopez <i>et al.</i> ⁽¹²⁾
Austria	Mineral water	3.7–211 mBq l ⁻¹	Wallner <i>et al.</i> ⁽¹³⁾
Brazil	Groundwater aquifer	0.01–1.5 Bq l ⁻¹	Bonotto <i>et al.</i> ⁽¹⁴⁾
Milano, Italy	Tap water	0.4–7.2 mBq l ⁻¹	Rusconi <i>et al.</i> ⁽¹⁵⁾
Nigeria	Mineral water	2.22 – 15.5 Bq l ⁻¹	Ajayi <i>et al.</i> ⁽¹⁶⁾
Italy	Bottled mineral water	< 10–52.5 mBq l ⁻¹	Desideri <i>et al.</i> ⁽¹⁷⁾
Kutahya, Turkey	Tap and spring waters	< 0.02–0.7+0.2 Bq l ⁻¹	Sahin <i>et al.</i> ⁽¹⁸⁾
Zahedan, Iran	Springs, wells and river	< 2-3 mBq l ⁻¹ (well water) 5 mBq l ⁻¹ (river) < 2 mBq l ⁻¹ (spring water)	Hosseini ⁽¹⁹⁾
Turkey	Mineral water	< 74 – 625 mBq l ⁻¹	Tabar <i>et al.</i> ⁽²⁰⁾
This work	Tap, spring and well waters	< 2- 38.2 m Bq l ⁻¹	2014

CONCLUSION

Natural exposures, such as gross- α , gross- β and ²²⁶Ra radionuclide were measured in the samples of Guilan province water supplies and springs. The results showed that the gross- α , gross- β and ²²⁶Ra concentration in all water samples were lower than WHO and EPA recommendations considered safe for consumption in study area.

ACKNOWLEDGMENT

The authors would like to thank the personnel of the National Radiation Protection Institute (NRPI) of Iran and Notional Geoscience Database of Iran (NGDI).

Conflict of interest: Declared none.

REFERENCES

1. Dragovic SD, Jankovic-Mandic LJJ, Dragovic RM, Cordevi MM, Cokic MM (2012) Spatial distribution of the ²²⁶Ra activity concentrations in well and spring waters in Serbia and their relation to geological formations. *Journal of*

Geochemical Exploration, **112**: 206-211.

2. EPA (Environmental Protection Agency) (2011) Radium in water. Illinois Department of Public Health, Division of Environmental Health, **217**: 782-5830.
3. EPA (Environmental Protection Agency) (1999) Radium-226 in drinking water radon emanation technique. *Method 903*, **1**: 38 – 48.
4. Mathieu G, Biscayne P, Lupton R, Hammond DE (1988) System for measurement of ²²²Rn at low level in natural waters. *Health Physics*, **55**: 989-992.
5. Hamlat MS, Kadi H, Djeflal S, Brahimi H (2003) Radon concentrations in Algerian oil and gas industry. *Applied Radiation and Isotopes*, **58**: 125-130.
6. Papastefanou C (2007) Measuring radon in soil gas and groundwater: a review. *Annals of Geophysics*, **50**: 569-578.
7. Smith CG, Cable JE, Martin J, Moutusi R (2008) Evaluating the source and seasonality of submarine groundwater discharge using a radon-222 pore water transport model. *Earth and Planetary Science Letters*, **273**: 312-322.
8. UNSCEAR (2000) Exposures from natural sources. Report to General Assembly, Annex B.
9. Porntepasemsan B and Srisuksawad K (2008) Assessment of Ra-226 age dependent dose from water intake. *Appl Radiat Isot*, **66** (11): 1654-1656.
10. WHO (World Health Organization) (2004) Guidelines for third edition recommendations drinking-water quality. *Vol.1, Geneva*.
11. Binesh A, Mohammadi S, Mowlavi AA, Parvaresh P (2011) Measuring radon and radium concentrations in 120 samples of drinking water sources, springs and rivers of Shandiz, Zoshk and Abrdeh regions. *Environ Res J*, **5**: 1–5.
12. Lopez MG and Sanchez AM (2008) Present status of Rn-222 in groundwater in Extremadura. *J Environ Radioact*,

Int. J. Radiat. Res., Vol. 14 No. 4, October 2016

- 99: 1539–1543.**
13. Wallner G, Wagner R, Katzlberger C (2008) Natural radionuclides in Austrian mineral water and their sequential measurement by fast methods. *J Environ Radioact*, **99**: 1090–1094.
 14. Bonotto DM and Bueno TO (2008) The natural radioactivity in Guarani aquifer groundwater, Brazil. *Appl Radiat Isot*, **66**: 1507–1522.
 15. Rusconi R, Forte M, Badalamenti P, Bellinzona S, Gallini R, Maltese S, Romeo C, Sgorbati G (2004) The monitoring of tap waters in Milano: planning methods and results. *Radiat Prot Dosim*, **111 (4)**: 73–376.
 16. Ajayi OS and Adesida G (2009) Radioactivity in some sachet drinking water samples produced in Nigeria. *Iran J Radiat Res*, **7 (3)**: 151-158.
 17. Desideri D, Meli MA, Feduzi L, Roselli C, Rongoni A, Saetta D (2007) ^{238}U , ^{234}U , ^{226}Ra , ^{210}Po concentrations of bottled mineral waters in Italy and their dose contribution. *J Environ Radioact*, **94**: 86–97.
 18. Sahin L, Çetinkaya H, Murat M, Ichedef M (2013) Determination of radon and radium concentrations in drinking water samples around the city of Kutahya. *Radiat Prot Dosim*, **155(4)**: 474–482.
 19. Hosseini SA (2007) Radioactivity monitoring in drinking water of Zahedan, Iran. *Iran J Radiat Res*, **5(2)**: 97-100.
 20. Tabar E and Yakut H (2014) Determination of ^{226}Ra concentration in bottled mineral water and assessment of effective doses, a survey in Turkey. *Int J Radiat Res*, **12(3)**: 193-201.